

Genome Sequences of Canine Parvovirus Type 2c Prevalent in Western Mexico

Elizondo Quiroga, D¹, De Los Santos Acuña, MA², Gutierrez Ortega, A¹, Galán Martínez, C², Pedroza Roldán, C^{2*}

1. *Unidad de Biotecnología Médica y Farmacéutica, Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco, A.C. Av. Normalistas No. 800. Col. Colinas de la Normal, CP. 44270, Guadalajara, Jalisco, México.*
2. *Departamento de Medicina Veterinaria, Centro Universitario de Ciencias Biológicas Y Agropecuarias, Cam. Ramón Padilla Sánchez 2100, Las Agujas, 44600 Zapopan, Jal. Universidad de Guadalajara, Zapopan, Jalisco, México.*

How to cite this article: ElizondoQuiroga D, DeLosSantosAcuña MA, GutierrezOrtega A, GalánMartínez C, PedrozaRoldán C. Genome Sequences of Canine Parvovirus Type 2c Prevalent in Western Mexico. *Archives of Razi Institute Journal*. 2024;79(2):387-394. DOI: 10.32592/ARI.2024.79.2.387



Copyright © 2023 by



Razi Vaccine & Serum Research Institute

ABSTRACT

Canine parvovirus type 2 (CPV-2) is one of the main etiologies of viral gastroenteritis in dogs across the globe. This disease is mainly characterized by the presence of diarrhea, abdominal pain, vomiting, anorexia, and dehydration. This virus is responsible for high mortality and morbidity rates in unvaccinated dogs and those younger than three months. The monitoring of viral variants in our region has demonstrated that in the last seven years, variant CPV-2c has been circulating exclusively, which is unusual if we consider that in the rest of the world, at least two variants co-circulate among dog populations. To the best of our knowledge, no studies in Mexico have reported genomic sequences of CPV-2, which are relevant for population comparisons at the genetic level. Therefore, the present study aimed to sequence genomes associated with CPV-2c. To meet this objective, rectal swab samples were collected from dogs with suspected CPV-2 infection. Five positive cases diagnosed by lateral flow testing and polymerase chain reaction were selected for viral genome sequencing. Comparative analyses illustrated that the obtained genome sequences were > 99% homologous to those reported for CPV-2 in the GenBank. On the other hand, 52 nucleotide mutations were identified in the *vp1/vp2* gene, out of which three impacted amino acid transition (T226S, F267Y, and A440T). Phylogenetic analysis of the *vp1/vp2* gene demonstrated that the five sequences clustered in a clade called "III", pertaining to sequences from USA and Uruguay. To our knowledge, this was the first report of genomic sequences associated with CPV-2 in Mexico, which is of great relevance for the epidemiological-molecular understanding and evolution of the virus.

Keywords: Canine parvovirus type 2c, Genome, Mexico

Article Info:

Received: 15 August 2023

Accepted: 30 September 2023

Published: 30 April 2024

Corresponding Author's E-Mail:

cesar.pedroza@academicos.udg.mx

1. Introduction

Canine parvovirus type 2 (CPV-2) is one of the main etiologic agents associated with gastroenteritis in unvaccinated dogs and those younger than three months. The gastroenteric picture is characterized by the presence of clinical signs, such as vomiting, diarrhea, abdominal pain, anorexia, and dehydration (1). CPV-2 is a non-enveloped virus comprising a positive-sense single-stranded DNA genome approximately 5.2 kb in size. The genome of CPV-2 has two open reading frames (ORFs), one of which encodes two nonstructural proteins (NS1 and NS2), and the other one encodes a protein precursor from which the structural proteins VP1 and VP2 are derived. The latter represents more than 80% of the total protein of the virus since it is responsible for the formation of the viral capsid (2). The NS1 protein is a 76-kDa nuclear phosphoprotein with multiple functions. One of its primary functions is viral replication, where the N-terminal portion of the protein recognizes the ORI site of the viral sequence in a strand-specific manner (3). On the other hand, the NS2 protein has been detected in the nucleus and cytoplasm, where it has been associated with viral replication (4). The VP1 protein contains amino acids in the N-terminal region responsible for transporting the viral capsid into the nucleus (5). The VP2 protein, which is in charge of cell tropism, contains antigenic sites highly relevant for antibody-mediated viral neutralization, and its amino acid composition is analyzed for epidemiological-molecular studies that allow the identification of preponderant viral variants in canid populations (6, 7). Soon after emergence, the original CPV-2 was replaced by two antigenic variants, namely CPV-2a and CPV-2b, which differ by an amino acid substitution (Asn to Asp) at position 426 of the VP2 protein. In 2000, a new antigenic variant that emerged in Italy was classified as CPV-2c, characterized by the presence of a glutamic acid (Glu) at position 426 (8). The three viral variants co-circulate worldwide, and their distribution and genetic diversity may vary depending on the regions studied (9). For instance, in Asian countries, there is a predominance of the CPV-2a and CPV-2b variants with a lower prevalence of the CPV-2c variant (10, 11). In European countries, there is no predominance of any of the variants; nonetheless, all three co-circulate in a balanced manner in different regions (12). On the contrary, in African countries, CPV-2b infection is predominant, with CPV-2a co-circulating to a lesser extent (13). In South American countries, such as Uruguay, Brazil, and Argentina, the predominance of CPV-2c variant circulation has been identified, compared to CPV-2a and CPV-2b (9). Nevertheless, in countries such as Ecuador, Colombia, and Peru, the predominance falls on the CPV-2a variant with a lower prevalence of

CPV-2b and CPV-2c (14, 15). Our group has previously reported an exclusive circulation of CPV-2c in the Western region of Mexico and possibly throughout the country, a phenomenon that could be considered unusual (16, 17). In addition, the analysis of partial sequences encoding the VP2 protein revealed that the CPV-2c circulating in our region is phylogenetically clustered; however, it is related to CPV-2c sequences reported in European and South American countries, suspecting that it was introduced to Mexico from these regions (17). Partial or complete genome sequences derived from CPV-2c have been previously reported in some countries; however, in Mexico, there are no studies on the characterization of the CPV-2 genome, which limits the knowledge of an infectious agent highly relevant to animal health. In light of the aforementioned issues, the present study aimed to sequence and analyze the genome of CPV-2c, which is prevalent in the Western region of Mexico.

2. Materials and Methods

2.1. Sample collection

Outbreaks of CPV-2 infection were identified in the municipalities of Guadalajara and Zapopan in the state of Jalisco, in western Mexico. A total of 22 of them underwent treatment at the Veterinary Hospital for Small Pets, affiliated with the University of Guadalajara, within two weeks. These dogs presented various signs, such as bloody diarrhea, vomiting, abdominal pain, anorexia, and dehydration. Rectal swabbing was performed, and the presence of CPV-2-associated antigen was confirmed by lateral flow CPV Ag Test Kit (Bionote, South Korea); rectal swab samples were frozen at -20°C .

2.2. Confirmation by polymerase chain reaction of the presence of CPV-2

DNA extraction and purification from fecal samples were performed using the GF-1 DNA extraction kit (Vivantis, Malaysia) following the manufacturer's instructions. DNA was eluted from columns with 50 μl of DNase and RNase-free water and stored at -20°C until use. Polymerase chain reaction (PCR) amplification of a partial segment within the *vp2* gene was performed with oligonucleotides described previously (18). The PCR reactions were performed in a final volume of 25 μl , with 2 μl of purified DNA, 2.5 μl of $10\times$ PCR buffer, 10 pmol of each primer, 0.5 μl of 10 mmol/l dNTPs mix, and 0.3 μl of Taq DNA polymerase (Vivantis, Malaysia) (5 U/ μl). Thermal cycler conditions were initial denaturation at 94°C for 5 min, followed by 30 cycles of denaturation at 94°C for 30 sec, annealing at 50°C for 1 min, extension at 72°C for 1 min, and a final extension at 72°C for 10 min. DNA from a commercially available, live attenuated vaccine was used as the positive control, while pure water

and DNA extracted from rectal swabs of healthy dogs were utilized as negative controls. The PCR products were separated in 1.5% agarose gels and stained with Sybr Safe (Vivantis, Malaysia). Samples were considered PCR-positive upon visual detection of the expected 1,042-bp amplicon after gel electrophoresis (16).

2.3. Genome sequencing and VP2 phylogenetic analysis

Four pairs of specific oligonucleotides were designed from the genomic sequence of CPV with accession number KF638400 previously reported in the GenBank database (NCBI) (Table 1). Serial Cloner software (version 2.6) was used to design and select the oligonucleotides. Of the 22 samples collected, 5 cases were selected for viral genome sequencing. The PCR reactions were performed in a final volume of 25 µl, with 3 µl of purified DNA, 2.5 µl of 10× PCR buffer, 10 pmol of each primer, 2 µl of 25 mM magnesium chloride, 0.5 µl of 10 mM dNTP mix, and 0.3 µl of high fidelity Taq DNA polymerase (Vivantis, Malaysia) (5 U/µl). The thermal cycling conditions included initial denaturation at 94°C for 5 min, followed by 25 cycles of denaturation at 94°C for 30 sec, alignment at 53.5°C for 30 sec, and extension at 72°C for 1 min. An elongation step was added at 72°C for 10 min, and the samples were preserved at -20°C. The PCR amplicons were separated in 1% agarose gels in the presence of Sybr-safe dye, and the products were visualized on an LED transilluminator. Bands with corresponding molecular weights (1100 bp, 1122 bp, 1384 bp, and 1329 bp) were obtained from the gel. The Wizard SV Gel and PCR Clean-up System kit (Promega, USA) was used to purify the PCR amplicons following the manufacturer's instructions. DNA was eluted from the columns with 50 µl of double distilled water, quantified, and its purity was determined in a Nanodrop spectrophotometer. The PCR amplicons were

submitted to the DNA Synthesis and Sequencing Unit of the Institute of Biotechnology, National Autonomous University of Mexico, for DNA sequencing using the Sanger method. The sequences were quality-checked and analyzed using the FinchTV (version 1.5) and UGENE (version 44) software packages. The identity of each amplicon was checked using the NCBI-BLAST algorithm. Genome assembly from the obtained sequences was performed using UGENE software (version 44) in conjunction with the read de novo assembly (CAP3). The genomes were verified in both sense and antisense sequences, and once finalized, their identities were determined using the NCBI-BLAST algorithm. The obtained genome sequences were submitted to the GenBank Database to obtain the accession numbers.

2.4. Alignments

To compare the sequences reported herein, CPV complete sequences were searched in GenBank. In this regard, 10 countries from various parts of the world were selected due to the sequence length (four from Asia, two from the Western Hemisphere, two from Europe, one from Africa, and one from Oceania). From each country, two sequences were used for the alignment, except for the USA, where an extra sequence was used to generate the comparison root, considering the proximity of the country to Mexico. The five sequences from Jalisco were cut to fit the size of sequences found at GenBank (4269 bp). Nucleotide of the entire sequences and deduced amino acid sequences of the VP2 protein were aligned and compared to sequences belonging to the selected countries using a progressive alignment algorithm in the CLC main Work Bench software package (Version 7.7.1.) (Quiagen, Denmark, <https://www.qiagenbioinformatics.com/>).

Table 1. Oligonucleotide description and used in this study.

Code	Oligonucleotides	Length of amplification (bp)	Coordinates*
F1F	5'-GAATGATAGGCGGTTTGTGT-3'	1 100	0-1100
F1R	5'-TCCTGGTTGTGCCATCATT-3'		
F2F	5'-CCTGAAGACTGGATGATGTTA-3'	1 122	1050-2172
F2R	5'-TGCCATCGTACCTTAATCCA-3'		
F3F	5'-CGTGGTCCGAAATAGAGGCA-3'	1 384	2078-3462
F3R	5'-TGTTCCCTGTAGCAAATTCATCACC-3'		
F4F	5'-ATATATACCATGGTACAGATC-3'	1 329	3359-4687
F4R	5'-TAACAACATACATTAATGGT-3'		

* The coordinates are based on the CPV-2 genome sequence with GenBank accession number: KF638400.

2.5. Phylogenetic trees

The evolutionary distances were computed using the Maximum Likelihood reconstruction with the Neighbor-joining method (Saitou and Nei, 1987) and the Jukes-Cantor nucleotide substitution model with rate variation. Phylogenetic tree construction was performed with 1,000 bootstrap replicates to support the information from the internal nodes, showing confidence values to demonstrate statistical support for each clade. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1,000 replicates) are displayed next to the branches. Only bootstrap values above 70% were indicated at the node. The analysis involved 26 complete sequences. All positions containing gaps and missing data were eliminated. There was a total of 4,269 positions in the final dataset. Evolutionary analyses were conducted in CLC main Work Bench software (version 7.7.1.).

3. Results

It is well known that VP2 is the protein that has higher variability, producing a wide host range and cross-reactions between variants. Therefore, the VP2 of our sequences was selected for genetic variation analysis. The five genome sequences presented almost identical homology compared to the US-17 sequence, presenting the highest homology rate in MX1-17 with 99.72%; the rest had homology rates of 99.46-99.67%. When comparing all sequences with US-17, 52 nucleotide mutations were identified in the *Vp2* gene, out of which 39 were synonymous and 13 involved amino acid substitutions (Table 2). When US-17 was compared with Mexico sequences (MX1-15, MX2-15, MX-16, MX1-17, and MX2-17), 12 and 3 synonymous and non-synonymous mutations were identified, respectively. One of the three identified mutations involves changing the amino acid threonine at position 226 to serine (T226S), which is present in 100% of the sequences compared to the US-17 sequence. The mutation at nucleotide 799 in VP2 resulted in an amino acid substitution at codon position 267 (F267Y). The amino acid substitution at residue 267 was present as a mutation in all sequences used and described from countries in Asia (IN-20, IN-16, KR1-17, CN1-19, TH1-16, TH2-16, CN2-19, and KR2-17), Africa (NG1-18 and NG2-18), and one from Uruguay (UY2-11). The mutation at codon position 440 (A440T) was observed in two Jalisco sequences (MX1-15 and MX2-17). This mutation was present in some of the sequences from all the regions we used in the analysis, such as France and Finland (FR-83, FR-84, FI-80, and FI-86), Australia (AU-82 and AU-85), Thailand, China, South Korea (TH1-16, TH2-16, CN2-19, and KR2-17), Uruguay, and USA (UY1-11 and US-15). To visualize the relationship of the CPV sequences, a phylogenetic tree was constructed based on the five full-length sequences

presented herein, and 21 sequences were retrieved from GenBank. The resulting phylogenetic tree demonstrated that all the Jalisco state sequences are grouped in a clade III, which is more related to sequences from the USA (US-15, US-17, and US-19) and one sequence from Uruguay (UY1-11) (Figure 1). Two more clades were generated separately, grouping the rest of the retrieved sequences. The clade I was divided into two clusters that grouped separately: Nigeria (NG1-18 and NG2-18), India (IN-16 and IN-20), China (CN1-19), and Uruguay (UY2-11) in cluster 1; and Thailand (KR1-17 and KR2-17), China (CN2-19), and Korea (KR2-17) sequences in cluster 2. Finally, clade II grouped Finland (FI-80 and FI-86) in one cluster, and France (FR-83 and FR-84) and Australia (AU-82 and AU-85) in the second cluster (Figure 1). The phylogenetic analysis considering the 21 complete CPV sequences illustrated that sequences MX2-15, MX-16, and MX1-17 were clustered together in a monophyletic clade as a branch of the root sequence from USA (US-17). Sequences MX1-15 and MXG-17 were in two different subgroups; the first was more related to US-15 (a strain from Ohio, USA) and the second pertained more closely to a strain from Uruguay (UY1-11).

4. Discussion

Previously, our group performed epidemiologic-molecular studies analyzing a fragment of the gene associated with the VP2 protein and found that the CPV-2c variant (426-Glu) has been circulating exclusively in our region and possibly in the country during the last seven years (16, 17, 19). To gain a deeper understanding of this virus, we sequenced five CPV-2c genomes. A homology of more than 99% with CPV-2 was obtained by comparing the sequences found here with others reported. The complete analysis of the VP2 protein-associated gene allowed the identification of 12 synonymous and 3 non-synonymous mutations (T226S, F267Y, and A440Y). In the present study, the MX2-15 genome sequence presented a replacement of Phe by Tyr at position 267. This finding was interesting since it is a mutation that had not been identified in our region and is very characteristic of Asian countries (20). This mutation has been reported to be present in CPV-2a VP2 in several groups, including China (21, 22), South Korea (23), and Nigeria (24). In addition, this mutation has occurred with high frequency in the CPV-2b viral variant identified in Asian countries where, in combination with other residues (25), it has been called "new CPV-2b" since approximately 92% of the analyzed strains present this mutation (20). The residue is in the greatest variable GH loop comprising amino acids 267-498 of the VP2 protein (26), which contains antigenic antibody recognition sites (20)




Figure 1. Phylogenetic analysis based on complete nucleotide sequences of CPV, retrieved from GenBank. The phylogenetic tree was constructed with Maximum Likelihood phylogeny using the neighbor-joining method with 1,000 bootstrap replicates in the CLC main Work Bench V. 7.7.1. software package. The scale bar indicates nucleotide substitutions per site.

Tracking this residue in future research in our region is of great importance. It is possible that viral variants with this mutation spread more frequently than others. The alanine-to-threonine substitution at position 440 was identified in all five genomes described in this study. In our previous analysis, we observed this substitution in 48% (14/29) of sequences (17), and Faz et al. reported identification in approximately 50% of sequences (19). This substitution is not limited to the CPV-2c viral variant but has also been observed in the CPV-2a viral variants in China (27) and South Africa (13), as well as in CPV-2b in Brazil (28). The relevance of this substitution has not been understood so far. Nonetheless, this residue is at the top of the threefold spike, a region characterized by high antigenicity (6). Therefore, it could be assumed that this residue could be receiving selection pressures due to vaccination, as previously reported (21,29). The phylogenetic analyses performed in the present study confirmed our previous findings, according to which the phylogenetic relationship of circulating viruses in our region was highly similar to those reported in such countries as the United States and Uruguay (17). In agreement with the analyses demonstrated in the present report and those we have previously performed (16, 17), Grecco et al. pointed out that European viral clades have

spread in South America (30), allowing us to suggest that the Mexican clades have a South American origin. In conclusion, in the present study, we reported the first partial genomes of CVP-2c in Mexico, a variant that has been circulating exclusively for at least seven years. Comparative analyses with other CPV-2 genomes have demonstrated that they are > 99% homologous, and although 52 nucleotide mutations have been identified in the *VPI/VP2* gene, only three can affect the amino acid transition. Although it is still challenging to determine why only the CPV-2c variant circulates exclusively in Mexico, we can suggest that this variant is the product of viral dispersal from South America.

Acknowledgment

The authors wish to thank the University of Guadalajara, which provided partial funding under the Program to Support the Improvement of Production Conditions of SNI Members and SNCA-PROSNI 2022.

Authors' Contribution

CPR and MADA conceived the study. MADA, DEQ, CGM, and AGO performed experiments. CPR and DEQ interpreted data. CPR and DEQ wrote the manuscript. All authors read and approved the final manuscript.

Ethics

The study was approved by the Biosafety, Research, and Ethics Committees of the Veterinary Hospital of the University of Guadalajara. The pets' owners were informed about the purpose and procedures of the study and conferred consent for taking a rectal swab sample from the dogs. The study followed Federal Regulations, such as NOM-087-SEMARNAT-SSA1-2002, NOM-033-ZOO-1995, and NOM-062-ZOO-1999.

Conflict of Interest

The author(s) declares no potential conflicts of interest with respect to the research, authorship, and publication of this article.

Data Availability

The data that support the findings of this study are available on request from the corresponding author.

References

- Iris K, Leontides LS, Mylonakis ME, Adamama-Moraitou K, Rallis T, Koutinas AF. Factors affecting the occurrence, duration of hospitalization and final outcome in canine parvovirus infection. *Res Vet Sci.* 2010;89(2):174-8.
- Yuan W, Parrish CR. Canine parvovirus capsid assembly and differences in mammalian and insect cells. *Virology.* 2001;279(2):546-57.
- Niskanen EA, Ihalainen TO, Kalliolinna O, Hakkinen MM, Vihinen-Ranta M. Effect of ATP binding and hydrolysis on dynamics of canine parvovirus NS1. *J Virol.* 2010;84(10):5391-403.
- Wang D, Yuan W, Davis I, Parrish CR. Nonstructural protein-2 and the replication of canine parvovirus. *Virology.* 1998;240(2):273-81.
- Vihinen-Ranta M, Wang D, Weichert WS, Parrish CR. The VP1 N-terminal sequence of canine parvovirus affects nuclear transport of capsids and efficient cell infection. *J Virol.* 2002;76(4):1884-91.
- Allison AB, Organtini LJ, Zhang S, Hafenstein SL, Holmes EC, Parrish CR. Single Mutations in the VP2 300 Loop Region of the Three-Fold Spike of the Carnivore Parvovirus Capsid Can Determine Host Range. *J Virol.* 2016;90(2):753-67.
- Langeveld JP, Casal JI, Vela C, Dalsgaard K, Smale SH, Puijk WC, et al. B-cell epitopes of canine parvovirus: distribution on the primary structure and exposure on the viral surface. *J Virol.* 1993;67(2):765-72.
- Buonavoglia C, Martella V, Pratelli A, Tempesta M, Cavalli A, Buonavoglia D, et al. Evidence for evolution of canine parvovirus type 2 in Italy. *J Gen Virol.* 2001;82(Pt 12):3021-5.
- Miranda C, Thompson G. Canine parvovirus: the worldwide occurrence of antigenic variants. *J Gen Virol.* 2016;97(9):2043-57.
- Nandi S, Kumar M. Canine parvovirus: current perspective. *Indian J Virol.* 2010;21(1):31-44.
- Saei HD, Javadi S, Akbari S, Hadian N, Zarza E. Molecular Characterization of canine parvovirus (CPV) antigenic variants from healthy and diarrheic dogs in Urmia region, Iran. *Iranian Journal of Veterinary Medicine.* 2017;11:9-19.
- Decaro N, Desario C, Addie DD, Martella V, Vieira MJ, Elia G, et al. The study molecular epidemiology of canine parvovirus, Europe. *Emerg Infect Dis.* 2007;13(8):1222-4.
- Dogonyaro BB, Bosman AM, Sibeko KP, Venter EH, van Vuuren M. Genetic analysis of the VP2-encoding gene of canine parvovirus strains from Africa. *Vet Microbiol.* 2013;165(3-4):460-5.
- la Torre D, Mafla E, Puga B, Erazo L, Astolfi-Ferreira C, Ferreira AP. Molecular characterization of canine parvovirus variants (CPV-2a, CPV-2b, and CPV-2c) based on the VP2 gene in affected domestic dogs in Ecuador. *Vet World.* 2018;11(4):480-7.
- Duque-Garcia Y, Echeverri-Zuluaga M, Trejos-Suarez J, Ruiz-Saenz J. Prevalence and molecular epidemiology of Canine parvovirus 2 in diarrheic dogs in Colombia, South America: A possible new CPV-2a is emerging? *Vet Microbiol.* 2017;201:56-61.
- Pedroza-Roldan C, Paez-Magallan V, Charles-Nino C, Elizondo-Quiroga D, De Cervantes-Mireles RL, Lopez-Amezcuca MA. Genotyping of Canine parvovirus in western Mexico. *J Vet Diagn Invest.* 2015;27(1):107-11.
- Pedroza-Roldan C, Hernandez-Almaraz MA, Elizondo-Quiroga D, Gutierrez-Ortega A, Acosta-Monroy CM, Charles-Nino C, et al. Exclusive circulation of canine parvovirus type 2c in the Guadalajara metropolitan area in western Mexico: a five-year study. *Arch Virol.* 2022.
- Perez R, Francia L, Romero V, Maya L, Lopez I, Hernandez M. First detection of canine parvovirus type 2c in South America. *Vet Microbiol.* 2007;124(1-2):147-52.
- Faz M, Martinez JS, Gomez LB, Quijano-Hernandez I, Fajardo R, Del Angel-Caraza J. Origin and genetic diversity of canine parvovirus 2c circulating in Mexico. *Arch Virol.* 2019;164(2):371-9.
- Wang H, Jin H, Li Q, Zhao G, Cheng N, Feng N, et al. Isolation and sequence analysis of the complete NS1 and VP2 genes of canine parvovirus from domestic dogs in 2013 and 2014 in China. *Arch Virol.* 2016;161(2):385-93.
- Wu H, Li X, Wang L, Liu Y, Tian K. Molecular epidemiological survey of canine parvovirus in domestic dogs in four provinces, China. *Virusdisease.* 2018;29(1):113-7.
- Li C, Tang J, Chen Z, Li Q, Huang Z, Wang Q, et al. Genetic characterization of the complete genome of a mutant canine parvovirus isolated in China. *Arch Virol.* 2018;163(2):521-5.
- Xu J, Guo HC, Wei YQ, Shu L, Wang J, Li JS, et al. Phylogenetic analysis of canine parvovirus isolates from

- Sichuan and Gansu provinces of China in 2011. *Transbound Emerg Dis.* 2015;62(1):91-5.
24. Ndiana LA, Odaibo GN, Olaleye DO. Molecular characterization of canine parvovirus from domestic dogs in Nigeria: Introduction and spread of a CPV-2c mutant and replacement of older CPV-2a by the "new CPV-2a" strain. *Virusdisease.* 2021;32(2):361-8.
 25. Nikbakhat G, Shahram S, Mohyedidini S. Detection of a new canine parvovirus mutant in Iran. *Iranian Journal of Veterinary Medicine.* 2018;12(1):7.
 26. Battilani M, Ciulli S, Tisato E, Prosperi S. Genetic analysis of canine parvovirus isolates (CPV-2) from dogs in Italy. *Virus Res.* 2002;83(1-2):149-57.
 27. Zhong Z, Liang L, Zhao J, Xu X, Cao X, Liu X, et al. First isolation of new canine parvovirus 2a from Tibetan mastiff and global analysis of the full-length VP2 gene of canine parvoviruses 2 in China. *Int J Mol Sci.* 2014;15(7):12166-87.
 28. Castro TX, Costa EM, Leite JP, Labarthe NV, Cubel Garcia RC. Partial VP2 sequencing of canine parvovirus (CPV) strains circulating in the state of Rio de Janeiro, Brazil: detection of the new variant CPV-2c. *Braz J Microbiol.* 2010;41(4):1093-8.
 29. Zhou P, Zeng W, Zhang X, Li S. The genetic evolution of canine parvovirus - A new perspective. *PloS one.* 2017;12(3):e0175035.
 30. Grecco S, Iraola G, Decaro N, Alfieri A, Alfieri A, Gallo Calderon M, et al. Inter- and intracontinental migrations and local differentiation have shaped the contemporary epidemiological landscape of canine parvovirus in South America. *Virus Evol.* 2018;4(1):vey011.