



Research Paper

Co-occurrence of *sea*, *sec*, and *tst* Enterotoxin Genes in
Staphylococcus aureus isolates From Clinical SourcesAmeneh Sadat Sadeghi¹, Ebrahim Babapour^{1*}, Majid Taati Moghadam², Reza Mirnejad³

1. Department of Microbiology, Ka.C., Islamic Azad University, Karaj, Iran.

2. Department of Microbiology, Faculty of Medicine, Guilan University of Medical Sciences, Rasht, Iran.

3. Molecular Biology Research Center, Systems Biology and Poisonings Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran.

**How to cite this article** Sadeghi AS, Babapour E, Taati Moghadam M, Mirnejad R. Co-occurrence of *sea*, *sec*, and *tst* Enterotoxin Genes in *Staphylococcus aureus* isolates From Clinical Sources. *Archives of Razi Institute Journal*. 2026; 81(1):33-40. <https://doi.org/10.32598/ARI.81.1.3362>**doi** <https://doi.org/10.32598/ARI.81.1.3362>

Article info:

Received: 29 Oct 2025

Accepted: 10 Dec 2025

Published: 01 Jan 2026

Keywords:

Antibiotic resistance,
Staphylococcus aureus,
Staphylococcal enterotoxins,
Toxic shock syndrome toxin
(TSST)

ABSTRACT

Introduction: *Staphylococcus aureus* is a Gram-positive bacterium that can cause various diseases in specific conditions by secreting various toxins. Enterotoxins and toxins toxic shock syndrome toxin (TSST) play a major role in pathogenesis. Enterotoxins and TSST are pyrogenic super antigens that react with the MHC II molecule. The aim of this study was to investigate the frequency of the *sea*, *sec*, and *tst* genes in *S. aureus* isolated from clinical sources.**Materials & Methods:** This study was performed on 100 *S. aureus* isolates from hospitals in Karaj, which were finally identified using biochemical methods. Antibiotics susceptibility testing was performed by disk diffusion on agar, and the multiplex polymerase chain reaction (PCR) method was used to identify *sea*, *sec*, and *tst* genes.**Results:** The highest resistance was observed to penicillin (92%), while the lowest resistance was observed to vancomycin (0%), and 48 isolates (48%) were identified as multi-drug resistant (MDR). Although 86 isolates (86%) had at least one of the analyzed genes, only one (1%) isolate showed the presence of all three *sea*, *sec*, and *tst* enterotoxin genes, and 36% of isolates had the *sea* and *tst* genes. Among the 86 isolates, 79% contained the *sea* gene, 5% contained the *sec* gene, and 43% had the *tst* gene. Statistical analysis revealed a significant correlation between the presence of the *tst* gene and MDR isolates.**Conclusion:** The presence of relevant genes in clinical isolates should be considered in disease control and management due to the importance of *S. aureus* enterotoxins and TSST genes and their role in the development and exacerbation of staphylococcal diseases. Additionally, the high prevalence of antibiotic-resistant isolates limits antibiotic treatment.

* Corresponding Author:

Ebrahim Babapour, Assistant Professor.

Address: Department of Microbiology, Faculty of Medicine, Guilan University of Medical Sciences, Rasht, Iran.

Tel: +98 (912) 1483410

E-mail: e.babapour@kiaui.ac.ir; e_babapour@yahoo.comCopyright © 2026 The Author(s);
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>).
Noncommercial uses of the work are permitted, provided the original work is properly cited.

1. Introduction

S*taphylococcus aureus* is a common pathogen that can inhabit various parts of the body and cause a variety of infections, such as skin and tissue infections, food poisoning, hospital-acquired infections, pneumonia, septic arthritis, endocarditis, osteomyelitis, foreign body infections, and sepsis [1, 2]. *S. aureus* has multiple virulence factors that contribute to its pathogenicity and bacterial colonization [3]. These virulence factors include drug resistance, enterotoxins, and toxic shock syndrome toxin (TSST). Enterotoxins and TSST are pyrogenic super antigens that react with the MHC II molecule, causing T lymphocytes to proliferate extensively and leading to damage through the release of high levels of cytokines. *S. aureus* has been identified as having more than 23 types of enterotoxins, which contribute to gastrointestinal poisoning and gastroenteritis [4-6]. The majority of *S. aureus* strains found in patients with toxic shock syndrome produce a harmful toxin called TSST-1, which can cause failure of vital organs and is often fatal [7-9]. In addition, antibiotic resistance is a significant problem in managing various hospital infections. It not only causes treatment failure in some cases but also increases hospitalization time and treatment costs [10-14]. Infections caused by *S. aureus* are becoming increasingly difficult to treat due to the widespread circulation and emergence of drug-resistant strains. Toxic shock syndrome is often treated with clindamycin and vancomycin. However, the excessive use and inappropriate prescription of antibiotics have led to an increase in antibiotic resistance, further complicating the treatment of toxic shock syndrome [7, 15]. Furthermore, there is a lack of research on human samples in Iran, as most of the existing studies have focused on food and animal sources. Therefore, further research is needed to investigate the frequency of the *sea*, *sec*, and *tst* genes in *S. aureus* isolated from clinical sources, to better understand the health risks that patients face.

2. Materials and methods

2.1. Bacterial samples and identification

The study was conducted in 2021 on 100 *S. aureus* isolates collected from different samples of patients and outpatients from Karaj City hospitals, including wound, blood, urine, sputum, nasal, and pharyngeal swabs. The samples were identified using specific culture media and various biochemical tests in a microbiology laboratory. All identified isolates were inoculated in nutrient broth containing 20% glycerol and stored at -20 °C for further experiments [16, 17].

2.2. Antibiotics susceptibility test

To determine antibiotic sensitivity, all isolates were tested using the agar disk diffusion method on Muller-Hinton agar medium. The testing was performed with 12 antibiotic disks obtained from Padtan Teb Co., including: oxacillin (1 µg), vancomycin (30 µg), ceftaxime (30 µg), trimethoprim-sulfamethoxazole (1.25/23.75 µg), ciprofloxacin (5 µg), erythromycin (15 µg), clindamycin (2 µg), ceftazidime (30 µg), gentamicin (10 µg), tetracycline (30 µg), penicillin (10 U), and chloramphenicol (30 µg). The results were reported as sensitive, intermediate, and resistant based on the inhibitory zone. *S. aureus* strain ATCC 25923 was used as a positive control. Isolates were classified as multidrug-resistant (MDR) if they were resistant to at least one antibiotic from three different antibiotic families, based on the results of the antibiotic sensitivity test [18, 19].

2.3. DNA extraction

Initially, we isolated *S. aureus* strains and extracted DNA using the BetaPrep Genomic DNA Extraction Kit from BETA BAYERN, Germany. The quantity and quality of the extracted DNA were assessed using optical density (OD) 260/280 ratios and agarose gel (1.5%) electrophoresis. The extracted DNA was then preserved at -20 °C for future use.

2.4. Triplex PCR reaction

The triplex PCR reaction was used to confirm the presence of the analyzed genes in the studied isolates. Specific primers (Table 1) were used to determine if the isolates had the *sea*, *sec*, and *tst* genes. The reaction mixture contained a final volume of 30 µL, consisting of 15 µL of Amplicon's Master Mix (which includes Maxer Mix 1X, Tris-HCl 0.5 M, MgCl₂ 2 mM, dNTPs 1.6 mM, Taq 0.04 Units/µL, and 0.5 µL of water), 1 µL (0.2 µM) of forward and reverse primer for each gene, 2 µL (20 ng) of template DNA, and 7 µL of double-distilled sterile distilled water. The genes were amplified using a thermal cycler (Applied Biosystem) under the following conditions: an initial denaturation at 96 °C for 5 minutes, followed by 35 cycles consisting of denaturation at 94 °C for 1 minute, annealing at 57 °C for 1 minute, and extension at 72 °C for 1 minute. After the final amplification, the temperature was kept at 72 °C for 10 minutes [20]. A negative control was used, where all reagents were used except for the template DNA. For the positive control, the standard strains of *S. aureus* ATCC 13565, ATCC 19095, and ATCC 25923 were used for *sea*, *sec*, and *tst* genes, respectively. The Triplex PCR reaction product




Figure 1. The frequency of *S. aureus* isolates in various clinical samples

was run on a 2% agarose gel with a 100 bp ladder and checked under ultraviolet light with a Gel document device [21, 22].

2.5. Statistical analyses

After that, we analyzed the results using Microsoft Excel 2010 and SPSS software, version 16. We used Cramer's V and chi-square test, and set the significance level at $P \leq 0.05$.

3. Results

3.1. Bacterial samples

A total of 100 *S. aureus* isolates were collected from clinical sources in Karaj, with 49 from women and 51 from men. The average age of the patients was 46.51 years. The isolates were obtained from various sample types, with the highest percentage from blood (70%) and the lowest from wounds (2%). Other samples were taken from urine, sputum, nasal, and pharyngeal swabs. The frequency of *S. aureus* isolates in various clinical samples is presented in Figure 1. Our study found a significant relationship between the sample type and the presence of the *sea* gene in the *S. aureus* isolates ($P < 0.05$).

3.2. Antibiotics susceptibility

The following sentences refer to Table 2, which contains details of the results related to the pattern of resistance and sensitivity to antibiotics. The results of the disk agar diffusion method for antibiotic sensitivity testing showed that the 92(92%) isolates were resistant to penicillin, while 82(82%) isolates were resistant to ceftazidime. Additionally, 47(47%) isolates were resistant to tetracycline, 43(43%) to erythromycin, and 38(38%) to ceftoxitin. Other antibiotics with less resis-

tance included ciprofloxacin (36%), oxacillin (34%), and clindamycin (33%). Moreover, 23(23%) isolates were resistant to gentamicin, 16(16%) isolates to trimethoprim-sulfamethoxazole, and only four (4%) isolates to chloramphenicol. Notably, no vancomycin-resistant isolates were observed in the study. The results of the antibiotic sensitivity test showed that 48(48%) out of the total isolates were identified as MDR.

3.3. Presence of enterotoxin genes

Out of 100 isolates studied, 14 did not have the *sea*, *sec*, and *tst* genes, while the remaining 86(86%) isolates had at least one of these genes. Among the 86 isolates, 79% had the *sea* gene, 5% had the *sec* gene, and 43% had the *tst* gene. Moreover, 4% had both the *sea* and *sec* genes, 36% had the *sea* and *tst* genes, and 2% had both *sec* and *tst* genes (Figure 2). Only one (1%) isolate showed the presence of all three *sea*, *sec*, and *tst* enterotoxin genes. Statistical analysis revealed a significant correlation between the presence of the *tst* gene and MDR isolates ($P < 0.05$). Moreover, the frequency of this gene was higher in MDR isolates. The gender of patients had a significant association with the presence of the *sec* and *tst* genes ($P < 0.05$). The *tst* gene was more prevalent in female patients, while the *sec* gene was more prevalent in male patients.

4. Discussion

S. aureus is a significant pathogen for humans and has been a leading cause of both community-acquired and hospital-acquired infections for several decades. Despite antibiotic treatment, this microorganism frequently causes severe complications in hospitalized patients, and its increasing drug resistance has made treatment challenging. Genetically, this bacterium possesses genes that

Table 1. Sequence of primers for *sea*, *sec* and *tst* genes in *S. aureus* isolates

Gene primer	Sequence (5' to 3')	Primer Length	Fragment Size (bp)	Ref.
<i>sea</i>	F:GGTTATCAATGTGCGGGTGG	20	102	[23]
	R:CGGCACTTTTTCTCTTCGG	20		
<i>sec</i>	F:AGATGAAGTAGTTGATGTGTATGG	24	451	[23]
	R:CACACTTTAGAATCAACCG	20		
<i>tst</i>	F:ACCCCTGTTCCCTTATCATC	20	326	[23]
	R:TTTTCAGTATTTGTAACGCC	20		

contribute to virulence, antibiotic resistance, and enterotoxin production, which can have dangerous effects on the host [7, 24]. Our study analyzed 100 *S. aureus* isolates and found that the highest resistance rate was observed for penicillin (92%), while the lowest resistance rate was observed for vancomycin (0.0%). The resistance pattern to other antibiotics was as follows: ceftazidime (82%), tetracycline (47%), erythromycin (43%), cefoxitin (38%), ciprofloxacin (36%), oxacillin (34%), clindamycin (33%), gentamicin (23%), and trimethoprim-sulfamethoxazole (16%). In this regards, Jafari-Sales et al. (2019) reported a penicillin resistance rate of 100% in *S. aureus*, which is consistent with the findings of another study [25]. In 2016, a study showed that no *S. aureus* isolates were resistant to vancomycin, which is consistent with the present study, but the highest percentage of antibiotic resistance

was found for clindamycin, oxacillin, and trimethoprim-sulfamethoxazole [26]. Reisi et al. (2014) reported that the highest percentage of antibiotic resistance was to penicillin and cefotaxime (100%), while the lowest was to vancomycin (0.5%), which is in accordance with our results [27]. Wu et al. (2023) also found that 100% of *S. aureus* isolates were resistant to penicillin, but no vancomycin-resistant isolates were found among the samples [28]. Another study found that the level of antibiotic resistance to penicillin among *S. aureus* isolates was 68.3%, which is similar to our results. However, this level of resistance was lower than what we reported [2]. In another study, 92.5% of *S. aureus* isolates were resistant to penicillin, and 10.5% were confirmed as vancomycin-intermediate *S. aureus* [29]. The results of these studies and our own demonstrate differences and similarities in the level of re-

Table 2. Results of antibiotic sensitivity in the tested isolates

Antibiotic	Sensitivity	Total Isolates (n=100)		
	Sensitive (%)	Resistance (%)	Intermediate (%)	
Oxacillin	66	34	-	
Cefoxitin	62	38	-	
Trimethoprim-sulfamethoxazole	83	16	1	
Clindamycin	67	33	-	
Ciprofloxacin	58	36	6	
Chloramphenicol	91	4	5	
Erythromycin	49	43	8	
Gentamicin	75	23	2	
Ceftazidime	1	82	17	
Penicillin	8	92	-	
Tetracycline	53	47	-	
Vancomycin	100	-	-	




Figure 2. Comparison of the frequency of different enterotoxin genes in 100 *S. aureus* isolates

sistance to different antibiotics, which can be attributed to various factors such as geographical region and the type and number of collected samples. However, what is certain is the increasing rate of resistance in these bacteria.

The molecular results of the presence of enterotoxin genes in *S. aureus* isolates showed that 79%, 5%, and 43% of the isolates carried *sea*, *sec*, and *tst* genes, respectively. It was found that 36% of the isolates had both *sea* and *tst* genes, 4% had *sea* and *sec* genes, and 2% had *sec* and *tst* genes. Additionally, 1% of the isolates showed a positive presence of all three genes. In this regard, Goli et al. (2018) conducted a study on 49 *S. aureus* isolates, of which 34.7% were positive for the *sea* gene [30]. Various researchers from different parts of the world have reported different frequencies of the *sea* gene in *S. aureus* isolates. Some studies similar to our results, such as those conducted by Katayoon et al. (2017) [31] and Rahimi et al. (2014) [32], have reported a high frequency of the *sea* gene in *S. aureus* isolates, with 86.2% and 100% of the isolates carrying the gene, respectively. On the other hand, other studies, including those by Asgarpoor et al. 2018 [1] and Nashev et al. (2007) [33], have reported a lower level of the *sea* gene in *S. aureus* isolates, with carrier rates ranging from 16% to 47.4%. The frequency of the *sec* gene has been investigated in various studies, revealing a range of *S. aureus* isolates that harbor the gene. For instance, Goli et al. (2018) reported that 10% of *S. aureus* isolates carried the gene [30]. Other studies, such as those conducted by Eshraghi et al. (2009) [3], and Saadati et al. [34], reported frequencies of the *sec* gene in *S. aureus* isolates of 1.6%, and 9.5%, respectively. The frequency of the *tst* gene, which is one of the important toxins in the virulence of *S. aureus*, has been investigated in different studies. Mohammad Jani et al. (2018) reported that the prevalence of the *tst* gene in *S. aureus* isolates was 43%, which is in agreement with our results [35]. Howev-

er, other studies have reported different frequencies of this gene in *S. aureus* isolates. For example, Ramazanzadeh et al. (2015) reported 81% [36], Parsonnet et al. (2008) reported 9% [37], and Becker et al. (2001) reported 18.2% [38]. After comparing the findings of various research studies with the outcomes of the current study, it is evident that some results are consistent while others vary. The discrepancies in the frequency of the genes analyzed can be attributed to diverse factors such as: geographical location, sample type, sample size, strain natural habitat, the overall health of the population being studied, the pattern of health behavior in clinical and community settings, and differences in the investigation methods and primers used in molecular studies.

5. Conclusion

Our study revealed high levels of enterotoxins and antibiotic resistance in *S. aureus* isolates, which can exacerbate hospital infections and contribute to the spread of antibiotic resistance. It is crucial to take action against the rise of *S. aureus* genes that produce enterotoxins and TSST in clinical sources. Treating infections caused by this bacterium can be challenging and lead to severe consequences. Therefore, prioritizing disease control and avoiding unnecessary antibiotic use is essential to prevent the spread of resistance.

Ethical Considerations

Compliance with ethical guidelines

The study was approved by the Research Ethics Committee of Karaj Branch, Islamic Azad University, Karaj, Iran (Code: IR.IAU.K.REC.1399.048). All ethical considerations were observed in the preparation of this manuscript.

Data availability

All data generated or analyzed during this study are included in the article.

Funding

This research did not receive any grant from funding agencies in the public, commercial, or non-profit sectors.

Authors' contributions

Conceptualization: Ebrahim Babapour and Reza Mirnejad; Methodology: Ebrahim Babapour; Investigation, formal analysis, and writing the original draft: Ameneh Sadat Sadeghi; Review and editing: Ebrahim Babapour, Reza Mirnejad and Majid Taati Moghadam; Project administration: Reza Mirnejad.

Conflict of interest

The authors declared no conflict of interest.

Acknowledgements

The authors of this article express their gratitude to the following individuals and institutions for their assistance in conducting the study: the staff of the clinical laboratory of Imam Ali and Shahid Rajaei Hospitals, the Razi Laboratory of Karaj, and the knowledgeable experts of the Microbiology and Molecular Laboratory of the Microbiology Department at [Islamic Azad University, Karaj Branch](#).

References

- [1] Asgarpour D, Bahrami M, Daneshamooz S, Ghasemi M. [Identification of *Staphylococcus aureus* Enterotoxin Genes of sea, seb and sec among Healthy Carriers in Ardabil City (Persian)]. *Iran J Med Microbiol.* 2018; 11(6):149-57. [\[Link\]](#)
- [2] Tabaei S, Kouhi Noghondar M, Mohammadzadeh M, Ataei L, Amel Jamehdar S. [Pattern of antibiotic resistance in methicillin-resistant *Staphylococcus aureus* (MRSA) strains isolated from clinical specimens: Imam Reza hospital in Mashhad (Persian)]. *Med J Mashhad Univ Med Sci.* 2016; 59(2):64-70. [\[DOI:10.22038/mjms.2016.7328\]](#)
- [3] Eshraghi S, Salehipour Z, Pourmand MR, Forushani AR, Salehi M, Amiri SA, et al. [Prevalence of *tst*, *entC*, *entA* and *entA/C* genes in *staphylococcus aureus* strains isolated from different foods (Persian)]. *Tehran Univ Med J.* 2009; 67(7):470-6. [\[Link\]](#)
- [4] Vasconcelos NG, Cunha MD. Staphylococcal enterotoxins: Molecular aspects and detection methods. *J Public Health Epidemiol.* 2010; 2(3):29-42. [\[Link\]](#)
- [5] Samidoust P, Delshad ME, Talemi RN, Mojtahedi K, Samidoust A, Jahangiri S, et al. Incidence, characteristics, and outcome of COVID-19 in patients on liver transplant program: a retrospective study in the north of Iran. *New Microbes New Infect.* 2021; 44:100935. [\[DOI:10.1016/j.nmni.2021.100935\]](#) [\[PMID\]](#)
- [6] Tarzjani SD, Kamalzadeh S, Moghadam MT, Ashoobi MT. Clinical challenge of co-infection of SARS-CoV-2 with influenza during the influenza circulation season: Suggestions for prevention. *Germs.* 2023; 13(2):188-91. [\[DOI:10.18683/germs.2023.1384\]](#) [\[PMID\]](#)
- [7] Serray B, Mohammed S, Hammouni A, Timinouni M, EL Azhari M. Frequency of the toxic shock syndrome toxin-1 gene in methicillin-resistant *staphylococcus aureus* isolated from pediatric hospital in Morocco. *Moroccan J Public Health.* 2022; 4(1):43-8. [\[Link\]](#)
- [8] Zuheir Majeed H, Mohamed Baqer Muhsin Y, Mohammed Sajet R. Investigating the effects of ginger on biofilm production from bacteria isolated from respiratory tract. *Iran J Vet Med.* 2024; 18(Special Issue):637-48. [\[DOI:10.32598/ijvm.18.specialissue.3\]](#)
- [9] Azizkhani M, Jafari F, Haghighi P, Dehghan M. Evaluating contamination level of raw and roasted nuts distributed in commercial markets in Mazandaran province, Iran. *Iran J Vet Med.* 2020; 14(2):167-76. [\[Link\]](#)
- [10] Moghadam MT, Chegini Z, Norouzi A, Dousari AS, Shariati A. Three-decade failure to the eradication of refractory *Helicobacter pylori* infection and recent efforts to eradicate the infection. *Curr Pharm Biotechnol.* 2021; 22(7):945-59. [\[DOI:10.2174/1389201021666200807110849\]](#) [\[PMID\]](#)
- [11] Taati Moghadam M, Hossieni Nave H, Mohebi S, Norouzi A. [The evaluation of connection between integrons class I and II and ESBL-producing and Non-ESBL *klebsiella pneumoniae* isolated from clinical samples, Kerman (Persian)]. *Iran J Med Microbiol.* 2016; 10(4):1-9. [\[Link\]](#)
- [12] Mohebi S, Golestani-Hotkani Z, Foulad-Pour M, Nazeri P, Mohseni F, Hashemizadeh Z, et al. Characterization of integrons, extended spectrum beta lactamases and genetic diversity among uropathogenic *Escherichia coli* isolates from Kerman, South East of Iran. *Iran J Microbiol.* 2023; 15(5):616-24. [\[DOI:10.18502/ijm.v15i5.13867\]](#) [\[PMID\]](#)
- [13] Rastegar S, Skurnik M, Tadjrobehkar O, Samareh A, Samareh-Najaf M, Lotfian Z, et al. Synergistic effects of bacteriophage cocktail and antibiotics combinations against extensively drug-resistant *Acinetobacter baumannii*. *BMC Infect Dis.* 2024; 24(1):1208. [\[DOI:10.1186/s12879-024-10081-0\]](#) [\[PMID\]](#)
- [14] Rastegar S, Skurnik M, Niaz H, Tadjrobehkar O, Samareh A, Hosseini-Nave H, et al. Isolation, characterization, and potential application of *Acinetobacter baumannii* phages against extensively drug-resistant strains. *Virus Genes.* 2024; 60(6):725-36. [\[DOI:10.1007/s11262-024-02103-5\]](#) [\[PMID\]](#)
- [15] Pines M. The Dangers of *Staphylococcus aureus* and Antimicrobial Resistance [MA thesis]. Florida: Lynn University; 2023. [\[Link\]](#)

- [16] Aldujaily AH, Hassoon KF, Salman DB, Bustani GS. Effect of rhyza stricta-synthesized copper nanoparticles on Staphylococcus aureus-infected wounds in rabbit. Iran J Vet Med. 2025; 19(2). [Link]
- [17] Selim AO, Abdel Salam MM, Abdallah Hassan RN, Mustafa GE, Mahdy ZA. The effect of some nano plant extract on bacteria producing biogenic amines isolated from minced meat. Iran J Vet Med. 2024; 18(4):510-6. [DOI:10.32598/ijvm.18.4.1005428]
- [18] Ashoobi MT, Asgary MR, Sarafi M, Fathalipour N, Pirooz A, Jafaryparvar Z, et al. Incidence rate and risk factors of surgical wound infection in general surgery patients: A cross-sectional study. Int Wound J. 2023; 20(7):2640-8. [DOI:10.1111/iwj.14137] [PMID]
- [19] Gholami M, Moshiri M, Ahanjan M, Salimi Chirani A, Hasannejad Bibalan M, Asadi A, et al. The diversity of class B and class D carbapenemases in clinical Acinetobacter baumannii isolates. Infez Med. 2018; 26(4):329-35. [PMID]
- [20] Moradi M, Norouzi A. [Prevalence of bla-CTX-M, bla-SHV, and bla-TEM genes and comparison of antibiotic resistance pattern in extended-spectrum β -lactamase producing and non-producing groups of Klebsiella pneumoniae Isolated from clinical samples in Kerman hospitals (Persian)]. J Adv BiomedSci. 2016; 6(1):120-8. [Link]
- [21] Hasannejad-Bibalan M, Sadeghi M, Hemmati H, Ashoobi MT, Yaghoubi T, Samadnia A, et al. A two-year study of microbiological characteristics of intravascular catheter-related bloodstream infections at Razi hospital, Iran. New Zealand J Med Lab Sci. 2021; 75(3):202-5. [Link]
- [22] Saeidi Y, Pournajaf A, Gholami M, Hasannejad-Bibalan M, Yaghoubi S, Khodabandeh M, et al. Determination of Helicobacter pylori virulence-associated genes in duodenal ulcer and gastric biopsies. Med J Islam Repub Iran. 2017; 31:95. [DOI:10.14196/mjiri.31.95] [PMID]
- [23] Mehrotra M, Wang G, Johnson WM. Multiplex PCR for detection of genes for S. aureus enterotoxins, exfoliative toxins, toxic shock syndrome toxin 1 and methicillin resistance. J Clin Microbiol. 2000; 38(3):1032-5. [DOI:10.1128/JCM.38.3.1032-1035.2000] [PMID]
- [24] Strich JR, Heil EL, Masur H. Considerations for empiric antimicrobial therapy in sepsis and septic shock in an era of antimicrobial resistance. J Infect Dis. 2020; 222(Suppl 2):S119-31. [DOI:10.1093/infdis/jiaa221] [PMID]
- [25] Jafari-Sales A, Jafari B. [Evaluation of the Prevalence of mec A Gene in staphylococcus aureus strains isolated from clinical specimens of hospitals and treatment centers (Persian)]. Pajouhan Sci J. 2019; 17(3):41-7. [DOI:10.52547/psj.17.3.41]
- [26] Abbasi S, Taei S, Zamanzad B. The prevalence of methicillin-resistant Staph. aureus strains producing enterotoxin A and B. Tehran Univ Med Sci J. 2016; 73(11):778-83. [Link]
- [27] Reisi M, Tajbakhsh E, Momtaz H. [Isolation and identification of antibiotic resistance genes in Staphylococcus aureus isolates from respiratory system infections in shahrekord, Iran (Persian)]. J Microb Biol. 2014; 3(10):97-106. [Link]
- [28] Wu X, Wang C, He L, Xu H, Jing C, Chen Y, et al. Clinical characteristics and antibiotic resistance profile of invasive MRSA infections in newborn inpatients: A retrospective multicenter study from China. BMC Pediatr. 2023; 23(1):264. [DOI:10.1186/s12887-023-04084-0] [PMID]
- [29] Nepal N, Mahara P, Subedi S, Rijal KR, Ghimire P, Banjara MR, et al. Genotypically confirmed vancomycin-resistant Staphylococcus aureus with vanB gene among clinical isolates in Kathmandu. Microbiol Insights. 2023; 16:11786361231183675. [DOI:10.1177/11786361231183675] [PMID]
- [30] Goli HR, Ahanjan M, Firouzi F, Nasrollahi M. [Prevalence of enterotoxin A and enterotoxin B Genes in Staphylococcus aureus strains isolated from hospitalized patients, medical personnel, and kitchen staff in two educational hospitals, Sari, Iran (Persian)]. J Mazandaran Univ Med Sci. 2018; 28(165):159-64. [Link]
- [31] Katayoon A, Naser H, Mehrooz D. Molecular Survey of the Frequency of sea and seb genes in staphylococcus aureus isolated from skin infections in Razi Hospital of Tehran. 2017. [Link]
- [32] Rahimi F, Bouzari M, Katouli M, Pourshafie M. Study of prevalence of Enterotoxin type A gene in Meticillin Resistant Staphylococcus aureus (MRSA) Hospitals in Tehran. Iran J Infect Dis. 2014; 19:59-68. [Link]
- [33] Nashev D, Toshkova K, Bizeva L, Akineden Ö, Lämmle C, Zschöck M. Distribution of enterotoxin genes among carriage-and infection-associated isolates of Staphylococcus aureus. Lett Appl Microbiol. 2007; 45(6):681-5. [DOI:10.1111/j.1472-765X.2007.02254.x] [PMID]
- [34] Saadati M, Babak B, Mehdi S, Mohammadjavad S. Detection of staphylococcus aureus enterotoxin type C isolated from healthy carriers by PCR. 2007; 117-27. [Link]
- [35] Mohammad Jani F, Amini K. [Detection of virulence (etA, etB and tst) and antibiotic resistance (mecA) genes in Staphylococcus Aureus strains isolated from clinical samples using multiplex-PCR method (Persian)]. Payavard Salamat. 2018; 11(6):610-7. [Link]
- [36] Ramazanzadeh R, Salimizand H, Shahbazi B, Narenji H. Prevalence of mecA gene of methicillin resistant Staphylococcus spp. Isolated from nosocomial infections and environmental specimens in Sanandaj Hospitals, Kurdistan, Iran. Res Mol Med. 2015; 3(3):38-42. [DOI:10.7508/rmm.2015.03.008]
- [37] Parsonnet J, Goering RV, Hansmann MA, Jones MB, Ohtagaki K, Davis CC, Totsuka K. Prevalence of toxic shock syndrome toxin 1 (TSST-1)-producing strains of Staphylococcus aureus and antibody to TSST-1 among healthy Japanese women. J Clin Microbiol. 2008; 46(8):2731-8. [DOI:10.1128/JCM.00228-08] [PMID]
- [38] Becker K, Haverkämper G, Von Eiff C, Roth R, Peters G. Survey of staphylococcal enterotoxin genes, exfoliative toxin genes, and toxic shock syndrome toxin 1 gene in non-Staphylococcus aureus species. European J Clin Microbiol Infect Dis. 2001; 20:407-9. [DOI:10.1007/s100960100523] [PMID]

This Page Intentionally Left Blank
