



Research Paper

Assessment of Oxidative Stress and Lipid Peroxidation
in Cats Infected With *Toxoplasma*Seyedeh Parastoo Yasini^{1*}, Mina Hedayati², Seyedeh Paniz Mobayen³, Marziyeh Kefayat⁴

1. Department of Clinical Sciences, Ka.C., Islamic Azad University, Karaj, Iran.

2. Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran.

3. Faculty of Veterinary Medicine, Ka.C., Islamic Azad University, Karaj, Iran.

4. Department of Parasitology, Ka.C., Islamic Azad University, Karaj, Iran.

Use your device to scan
and read the article online**How to cite this article** Yasini SP, Hedayati M, Mobayen SP, Kefayat M. Assessment of Oxidative Stress and Lipid Peroxidation in Cats Infected With *Toxoplasma*. *Archives of Razi Institute Journal*. 2026; 81(3):613-618. <https://doi.org/10.32598/ARI.81.3.3566> <https://doi.org/10.32598/ARI.81.3.3566>

Article info:

Received: 19 Dec 2025

Accepted: 26 Mar 2026

Published: 01 May 2026

Keywords:

Catalase (CAT), Glutathione peroxidase (GPx), Malondialdehyde (MDA), Superoxide dismutase (SOD), Total antioxidant capacity (TAC), *Toxoplasma gondii*

ABSTRACT

Introduction: Toxoplasmosis, caused by the intracellular protozoan *Toxoplasma gondii*, is a widespread zoonotic infection that frequently remains asymptomatic in humans and animals. During parasitic infections, oxidative stress can increase due to excessive free radical production, potentially contributing to disease pathogenesis. The present study investigates oxidative stress biomarkers in cats, the definitive hosts of *T. gondii*.

Materials & Methods: Blood samples were collected from 55 cats. Among them, 10 infected and 10 un-infected cats were selected based on serological screening using the modified agglutination test (MAT). Serum levels of malondialdehyde (MDA), superoxide dismutase (SOD), glutathione peroxidase (GPx), catalase (CAT), and total antioxidant capacity (TAC) were analyzed. Data were evaluated using independent t-tests, with statistical significance set at $P \leq 0.05$.

Results: Although serum MDA concentrations were elevated in infected cats ($10.68 \pm 1.07 \mu\text{M}$) compared to controls ($9.12 \pm 0.56 \mu\text{M}$), the difference was not statistically significant. Enzyme activities of SOD showed no significant difference. Infected cats had lower GPx ($128.66 \pm 35.62 \text{ U/mL}$ vs $235.66 \pm 74.13 \text{ U/mL}$), CAT ($12.19 \pm 1.03 \text{ U/mL}$ vs $13.12 \pm 2.60 \text{ U/mL}$), and TAC ($0.118 \pm 0.004 \text{ mM}$ vs $0.126 \pm 0.005 \text{ mM}$), but these reductions were not statistically significant ($P > 0.05$).

Conclusion: This study evaluated serum oxidative markers and lipid peroxidation in cats naturally infected with *T. gondii*. The lack of significant differences suggests that oxidative stress and lipid peroxidation induced by the infection may be localized within cells, consistent with the intracellular nature of *T. gondii*, an obligate intracellular protozoan. Moreover, the absence of overt clinical signs in infected cats might account for the minimal changes in systemic antioxidant parameters. Further research incorporating tissue-specific analyses and experimental infection models is necessary to more accurately elucidate the oxidative stress mechanisms associated with *T. gondii* infection.

* Corresponding Author:

Seyedeh Parastoo Yasini, Assistant Professor.

Address: Department of Clinical Sciences, Ka.C., Islamic Azad University, Karaj, Iran.

E-mail: pyasini@kiauo.ac.irCopyright © 2026 The Author(s);
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>).
Noncommercial uses of the work are permitted, provided the original work is properly cited.

1. Introduction

Toxoplasmosis is a globally prevalent zoonotic disease caused by the intracellular protozoan *Toxoplasma gondii*, which affects a broad range of warm-blooded animals, including humans, mammals, and birds [1, 2]. Felids, especially domestic cats, are the only known definitive hosts capable of shedding oocysts through their feces, making them key contributors to environmental contamination. Intermediate hosts, such as humans and livestock, become infected through ingestion of tissue cysts in undercooked meat, accidental contact with oocyst-contaminated sources, or congenital transmission [2, 3].

The parasite's life cycle includes three infective stages: tachyzoites, bradyzoites, and sporozoites. Sporulated oocysts represent a main source of infection for humans and other intermediate hosts [2, 3]. All cat breeds, regardless of age or sex, are susceptible to *T. gondii* infection. Oocysts excreted in cat feces contaminate soil, water, vegetables, and animal forage. It is estimated that approximately 1% of the cat population is actively shedding oocysts at any given time. The shedding period typically lasts between one and two weeks, which limits the utility of fecal testing as a reliable diagnostic method. Remarkably, under optimal environmental conditions such as moist soil and moderate temperatures, *T. gondii* oocysts can remain infectious for over a year, supported by their highly resistant outer wall that provides protection against most common disinfectants [4-6]. Consequently, environmental contamination with oocysts represents a significant and persistent source of infection for herbivores, omnivores, carnivores, and humans alike. Like other members of the Apicomplexa phylum, *T. gondii* is an obligate intracellular parasite. A critical host defense mechanism is the oxidative burst, in which immune cells rapidly produce reactive oxygen species (ROS) and reactive nitrogen species (RNS) in response to infection. To survive, the parasite must counteract oxidative damage within host cells. Oxidative stress occurs when ROS production surpasses the host's antioxidant defenses, leading to cellular damage. The antioxidant system comprises both enzymatic components such as superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), and glutathione reductase and non-enzymatic elements like reduced glutathione (GSH) and uric acid [7-9]. Given the relevance of oxidative stress in parasitic infections, biomarkers such as MDA, SOD, CAT, and GPx provide valuable insights into disease progression [9-11].

During infection, the host's immune response intensifies ROS and RNS production to eliminate pathogens. However, these reactive species may simultaneously damage host tissues, contributing to disease pathology [11-13]. Thus, oxidative imbalance plays a major role in the pathogenesis of toxoplasmosis in humans and animal hosts [14]. This study aims to assess serum oxidative stress and lipid peroxidation markers in cats naturally infected with *T. gondii*.

2. Materials and Methods

2.1. Sample collection

To obtain 10 confirmed *T. gondii*-positive samples, 55 blood samples were collected from cats displaying clinical signs such as lethargy and swollen lymph nodes. An additional 10 samples were obtained from healthy cats presented for routine health checks, with no apparent clinical symptoms. Approximately 5 mL of whole blood were collected from the cephalic vein of each animal using sterile syringes and transferred into serum-separation tubes pre-coated with clot activators. The samples were centrifuged at 3000 rpm for 10 minutes using a Hettich Rotina 380 centrifuge (Andreas Hettich GmbH & Co. KG, Germany) to separate the serum, which was then aliquoted into microtubes and stored at -80 °C until supplementary analysis. Seropositivity was determined using the modified agglutination test (MAT), with a titer of $\geq 1:20$ considered positive. From the symptomatic cats, only those with confirmed seropositivity ($n=10$) were included in the infected group. The un-infected group comprised 10 cats with undetectable antibody titers ($<1:20$).

2.2. Serological testing: MAT

Serological testing was performed to detect *T. gondii* infection by identifying anti-*Toxoplasma* antibodies in serum samples. The presence of antibodies was assessed using the MAT with microplates containing U-shaped wells. For the assay, 50 μ L of 0.2 M 2-mercaptoethanol prepared in phosphate-buffered saline (PBS) was added to each well, followed by the serum samples. Serial two-fold dilutions of the sera were prepared, ranging from 1:10 to 1:100. Subsequently, 50 μ L of *T. gondii* tachyzoite antigen, suspended in an alkaline buffer, was added to each well, introducing approximately 1,000,000 tachyzoites per well. The plates were gently shaken and incubated at 37 °C for 24 hours prior to interpretation.

Positive samples were identified by diffuse agglutination, where the precipitate did not settle at the bottom of the well. Negative samples were indicated by the formation of a button-shaped sediment at the bottom of the well. Positive and negative controls were included on each plate for validation. Serum dilutions of 1:20 or higher were considered seropositive [2].

2.3. Measurement of MDA concentration

MDA concentrations, representing lipid peroxidation, were quantified using a commercially available assay kit (ZellBio GmbH, Germany). In this colorimetric method, samples were acidified and reacted with thiobarbituric acid (TBA) under high-temperature conditions, forming a colored complex. The intensity of the resulting pink product correlates with MDA concentration and was measured spectrophotometrically at 535 nm using a spectrophotometer (Genway Scientific Ltd., UK).

2.4. Measuring the activity of antioxidant enzymes

2.4.1. SOD enzyme measurement

SOD activity was measured using a commercial assay kit (ZellBio GmbH, Germany). The assay is based on the conversion of superoxide radicals to molecular oxygen and hydrogen peroxide. The final chromogenic compound was detected at 420 nm using an ELISA microplate reader (BioTek ELx808, Agilent Technologies, USA).

2.4.2. GPx enzyme measurement

GPx activity was assessed with a commercial kit (ZellBio GmbH, Germany), measuring the absorbance of the reaction endpoint at 405 nm via an ELISA reader (BioTek ELx808, Agilent Technologies, USA).

2.4.3. Measurement of CAT enzyme

CAT activity was evaluated employing a commercial kit (ZellBio GmbH, Germany), with absorbance readings taken at 405 nm on an ELISA reader (BioTek ELx808, Agilent Technologies, USA).

2.4.4. Total antioxidant capacity measurement

TAC was evaluated with a commercial kit (ZellBio GmbH, Germany). Absorbance of the resultant product was read between 460–490 nm using an ELISA microplate reader (BioTek ELx808, Agilent Technologies, USA).

2.5. Statistical analysis

All results were expressed as Mean±SE. Data analysis was conducted using SPSS software (version 24). Independent samples t-tests were applied to compare the mean values between infected and un-infected groups. A $P \leq 0.05$ was considered statistically significant for all comparisons.

3. Results

The study identified 10 cats as positive for *T. gondii* infection, based on antibody titers $\geq 1:20$. Additionally, 10 cats with undetectable antibody titers, indicating no evidence of *T. gondii* infection, were included in the un-infected (control) group. The statistical data on CAT, SOD, and GPx enzyme activities, as well as TAC, are presented in Table 1. Although the activities of all enzymes were lower in infected cats compared to un-infected cats, the decrease was not statistically significant ($P > 0.05$).

There were no statistically significant differences between the groups ($P > 0.05$).

3.1. MDA level

Figure 1 illustrates the MDA concentrations in both infected and un-infected groups. Although MDA levels appeared elevated in the infected group relative to controls, statistical analysis indicated that the difference did not reach significance ($P > 0.05$).

4. Discussion

In recent years, the role of oxidative stress in the development and progression of parasitic infections has garnered increasing attention. Several investigations have demonstrated elevated levels of ROS following parasitic infections, suggesting that oxidative imbalance may contribute significantly to disease mechanisms. This study evaluated the status of oxidative stress and lipid peroxidation in cats naturally infected with *T. gondii*, comparing results with un-infected controls.

Our findings indicate that lipid peroxidation, as assessed by MDA levels, was higher in infected cats; however, the increase did not reach statistical significance. Similarly, SOD levels in infected cats did not show significant differences compared to healthy cats. While CAT, GPx, and TAC levels showed a downward trend in infected cats, these changes were also not statistically significant. These findings suggest that oxidative altera-

Table 1. Mean values $\pm$ SE for TAC, CAT, SOD, and GPx activities in infected and un-infected cats

Parameter	Mean $\pm$ SE	
	Infected	Un-infected
TAC (mM)	0.118 $\pm$ 0.004	0.126 $\pm$ 0.005
SOD (U/mL)	34.47 $\pm$ 1.32	30.82 $\pm$ 3.82
GPx (U/mL)	128.66 $\pm$ 35.62	235.66 $\pm$ 74.13
CAT (U/mL)	12.19 $\pm$ 1.03	13.12 $\pm$ 2.6

tions induced by *T. gondii* infection may be primarily localized within tissues rather than reflected in serum enzyme levels. It is possible that, during the acute phase of toxoplasmosis, ROS are intensively produced, and oxidative stress is induced in the tissues of infected animals as the host defense against the infection [15, 16].


Comparable studies have reported similar findings. For instance, Engin et al. noted that mice infected with *T. gondii* exhibited increased MDA concentrations in liver, brain, and spleen, while no significant elevation was detected in serum levels [17]. Bahrami et al. also documented a non-significant rise in serum lipid peroxidation markers in infected rats on the eighth day post-infection [18].

In contrast, Al-Kennany reported a significant increase in MDA levels in placental tissues of ewes infected with *T. gondii*, suggesting that localized oxidative stress may vary depending on the specific tissue affected [19]. Atmaca et al. demonstrated a significant rise in MDA levels in gerbils infected with *Toxoplasma*, attributing this to

excessive production of free radicals and oxidative stress following infection [20].

With respect to SOD activity, our findings align with prior reports indicating no significant alterations in serum SOD levels in *T. gondii* infection [19, 21]. Nevertheless, some studies have shown reduced SOD activity in infected animals [20]. The research of Nazarlou et al. on experimental *T. gondii* infection in male rats demonstrated a significant reduction in SOD activity by day 80 post-infection, suggesting that prolonged infection may have a more pronounced impact on antioxidant defense mechanisms [22].

In terms of GSH peroxidase and catalase activity, our results demonstrated a reduction in infected cats, although these changes were not statistically significant. These results are consistent with previous investigations by Al-Kennany and Machado et al., who documented decreased GPx activity in infected tissues [9, 19]. Nazarlou also identified a significant reduction in catalase levels in rats infected with *T. gondii* [22]. Mohammed et al. reported reduced GPx activity in the serum of seroposi-

**Figure 1.** Mean MDA concentration in infected and un-infected cats

tive pregnant women [23]. In contrast, Delavari et al. observed increased GPx in liver tissue at an early stage of infection, suggesting time-dependent and tissue-specific differences in enzyme responses [24].

In addition, Türkoglu et al. reported a significant increase in GPx activity in the liver tissue of infected rats, whereas no significant changes were reported in brain and kidney tissues [21]. The TAC trends observed in our study are consistent with findings by Mohammed and Bahrami, who reported no significant reduction in serum catalase levels in infected samples [18, 23]. However, Delavari et al. observed an increase in TAC in liver tissue on the eighth day post-infection, indicating that antioxidant responses may fluctuate depending on the phase of infection [24].

5. Conclusion

The present study suggests that *T. gondii* infection affects antioxidant enzyme activity and lipid peroxidation in cats. However, these changes were not statistically significant in serum samples. Given that *T. gondii* is an obligate intracellular protozoan, oxidative stress may be more pronounced at the tissue level rather than in systemic circulation, which could explain the lack of significant variations in serum antioxidant markers. Further research focusing on tissue-specific analysis and experimental infections is necessary to better understand the oxidative stress mechanisms associated with *Toxoplasma* infections and their potential impact on host pathophysiology.

Compliance with ethical guidelines

It is hereby asserted that all ethical standards have been observed in the preparation of the submitted article.

Funding

This research did not receive any grant from funding agencies in the public, commercial, or non-profit sectors.

Authors' contributions

Conceptualization and study design: Seyedeh Parastoo Yasini; Data acquisition: Mina Hedayati and Seyedeh Paniz Mobayen; Experiments and data interpretation: Seyedeh Parastoo Yasini, Mina Hedayati, and Marziyeh Kefayat; Statistical analysis: Seyedeh Parastoo Yasini and Seyedeh Paniz Mobayen; Writing the original draft: Seyedeh Parastoo Yasini and Mina Hedayati; Review and editing: Seyedeh Paniz Mobayen and Mina Hedayati.

Conflict of interest

The authors declared no conflict of interest.

Data availability

The data that support the findings of this study are available on request from the corresponding author.

Reference

- [1] Bogitsh BJ, Carter CE, Oeltmann TN. Human Parasitology. 5th ed. Cambridge: Academic Press; 2018. [Link]
- [2] Dubey JP. Toxoplasmosis of Animals and Humans. Boca Raton, FL: CRC Press; 2021. [DOI:10.1201/9781003199373]
- [3] Delgado IL, Zúquete S, Santos D, Basto AP, Leitão A, Nolasco S. The apicomplexan parasite *Toxoplasma gondii*. Encyclopedia . 2022; 2(1):189-211. [DOI:10.3390/encyclopedia2010012]
- [4] Boothroyd JC. What a Difference 30 Years Makes! A Perspective on Changes in Research Methodologies Used to Study *Toxoplasma gondii*. Methods Mol Biol. 2020; 2071:1-25. [DOI:10.1007/978-1-4939-9857-9_1] [PMID]
- [5] Usey MM, Huet D. Parasite powerhouse: a review of the *Toxoplasma gondii* mitochondrion. J Eukaryot Microbiol. 2022; 69(6):e12906. [DOI:10.1111/jeu.12906] [PMID]
- [6] Moradi F, Dashti N, Farahvash A, Baghaei Naeini F, Zarebavani M. Curcumin ameliorates chronic *Toxoplasma gondii* infection-induced affective disorders through modulation of proinflammatory cytokines and oxidative stress. Iran J Basic Med Sci. 2023; 26(4):461-7. [PMID]
- [7] Li TT, Zhao DY, Liang QL, Elsheikha HM, Wang M, Sun LX, et al. The antioxidant protein glutaredoxin 1 is essential for oxidative stress response and pathogenicity of *Toxoplasma gondii*. FASEB J. 2023; 37(6):e22932. [DOI:10.1096/fj.202201275R] [PMID]
- [8] Liu Q, Wang ZD, Huang SY, Zhu XQ. Diagnosis of toxoplasmosis and typing of *Toxoplasma gondii*. Parasites & Vectors. 2015; 8:292. [DOI:10.1186/s13071-015-0902-6] [PMID]
- [9] Machado VS, Bottari NB, Baldissera MD, Rech VC, Ianiski FR, Signor C, et al. Diphenyl diselenide supplementation in infected mice by *Toxoplasma gondii*: Protective effect on behavior, neuromodulation and oxidative stress caused by disease. Experimental Parasitology. 2016; 169:51-8. [DOI:10.1016/j.exppara.2016.07.006] [PMID]
- [10] Kiral F, Karagenc T, Pasa S, Yenisey C, Seyrek K. Dogs with Hepatozoon canis respond to the oxidative stress by increased production of glutathione and nitric oxide. Veterinary Parasitology. 2005; 131(1-2):15-21. [DOI:10.1016/j.vet-par.2005.04.017] [PMID]
- [11] Dubey JP, Lindsay DS. Neosporosis, toxoplasmosis, and sarcocystosis in ruminants. Vet Clin North Am Food Anim Pract. 2006; 22(3):645-71. [PMID]

- [12] Sturge CR, Yarovinsky F. Complex immune cell interplay in the gamma interferon response during *Toxoplasma gondii* infection. *Infect Immun*. 2014; 82(8):3090-7. [DOI:10.1128/IAI.01722-14] [PMID]
- [13] Rahimi MT, Daryani A, Sarvi S, Shokri A, Ahmadpour E, Mizani A, et al. Cats and *Toxoplasma gondii*: A systematic review and meta-analysis in Iran. *Onderstepoort J Vet Res*. 2015; 82(1):e1-e10. [DOI:10.4102/ojvr.v82i1.823] [PMID]
- [14] Bottari NB, Mendes RE, Baldissera MD, Bochi GV, Moresco RN, Leal ML, et al. Relation between iron metabolism and antioxidant enzymes and δ ALA D activity in rats experimentally infected by *Fasciola hepatica*. *Exp Parasitol*. 2016; 165:58-63. [DOI:10.1016/j.exppara.2016.03.012] [PMID]
- [15] Ibrahim NI, Nada SM, Ahmed M, Salama M, El Rahman AE. Overview about oxidative stress in toxoplasmosis. *Eur Chem Bull*. 2023; 12(1):307-21. [Link]
- [16] Zewczyk Golec K, Pawłowska M, Wesołowski R, Wróblewski M, Mila Kierzenkowska C. Oxidative stress as a possible target in the treatment of toxoplasmosis: Perspectives and ambiguities. *Int J Mol Sci*. 2021; 22(11):5705. [DOI:10.3390/ijms22115705] [PMID]
- [17] Engin AB, Dogruman AI F, Ercin U, Celebi B, Babur C, Bukan N. Oxidative stress and tryptophan degradation pattern of acute *Toxoplasma gondii* infection in mice. *Parasitol Res*. 2012; 111(4):1725-30. [DOI:10.1007/s00436-012-3015-6] [PMID]
- [18] Bahrami S, Shahriari A, Tavalla M, Azadmanesh S, Hamidinejat H. Blood levels of oxidant/antioxidant parameters in rats infected with *Toxoplasma gondii*. *Oxid Med Cell Longev*. 2016; 2016:8045969. [DOI:10.1155/2016/8045969] [PMID]
- [19] Al-Kennany ER. Oxygen free radicals released in placentae of ewes naturally infected with *Toxoplasma gondii*. *Al-Anbar J Vet Sci*. 2009; 2(2):1-6. [Link]
- [20] Atmaca N, Çınar M, Güner B, Kabakci R, Gazıyağcı AN, Atmaca HT, et al. Evaluation of oxidative stress, hematological and biochemical parameters during *Toxoplasma gondii* infection in gerbils. *Ankara Univ Vet Fak Derg*. 2015; 62(3):165-70. [DOI:10.1501/Vetfak_0000002675]
- [21] Türkoglu SA, Yaman K, Orallar H, Camsari C, Karabork S, Ayaz E. Acute toxoplasmosis and antioxidant levels in the liver, kidney and brain of rats. *Ann Parasitol*. 2018; 64(3):241-7. [PMID]
- [22] Hoseiny Asl Nazarlu Z, Matini M, Bahmanzadeh M, Foroughi-Parvar F. *Toxoplasma gondii*: A Possible Inducer of Oxidative Stress in Reproductive System of Male Rats. *Iran J Parasitol*. 2020; 15(4):521-9. [DOI:10.18502/ijpa.v15i4.4857] [PMID]
- [23] Mohammed MA, Rajab KI, Al Rawi KF, Al Darwesh MY. Evaluation some antioxidants and oxidative stress index in seropositive toxoplasmosis in pregnant women in Ramadi city of Iraq. *Sys Rev Pharm*. 2020; 11:701-5. [Link]
- [24] Delavari I, Khadjeh G, Bahrami S, Jalali SM, Esmaealzadeh S. Evaluation of Local tissue oxidative stress and its possible involvement in the pathogenesis of toxoplasmosis in rats experimentally infected with *Toxoplasma gondii*. *Trop Biomed*. 2017; 34(3):708-16. [PMID]