



## Research Paper

Utilization of *Solanum nigrum* L. (Black Nightshade) Fruit  
as a Phytobiotic in Laying QuailSyauqi Syah Dewan<sup>1</sup>, Widya Hermana<sup>2\*</sup>, Sumiati Sumiati<sup>2</sup>

1. Study Program of Nutrition and Feed Science, Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, IPB University, Bogor, Indonesia.

2. Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, IPB University, Bogor, Indonesia.



**How to cite this article** Dewan SS, Hermana W, Sumiati S. Utilization of *Solanum nigrum* L. (Black Nightshade) Fruit as a Phytobiotic in Laying Quail. *Archives of Razi Institute Journal*. 2026; 81(1):165-172. <https://doi.org/10.32598/ARI.81.1.3679>

<https://doi.org/10.32598/ARI.81.1.3679>

## Article info:

Received: 07 Aug 2025

Accepted: 23 Nov 2025

Published: 01 Jan 2026

## ABSTRACT

**Introduction:** Black nightshade (*Solanum nigrum* L.) is known to contain bioactive compounds with antioxidant properties. This study aimed to evaluate the effects of black nightshade fruit extract as a phytobiotic on the performance and egg quality of laying quails (*Coturnix-coturnix japonica*).

**Materials & Methods:** A total of 160 female quails, aged six weeks, were randomly assigned to four treatment groups with four replications, each containing ten birds. The treatments included a control group (T0) and three experimental groups receiving drinking water supplemented with 2 mL/L (T1), 4 mL/L (T2), and 6 mL/L (T3) of black nightshade extract. The parameters observed included feed intake, quail-day production, egg production, egg weight, feed conversion ratio (FCR), physical egg quality (egg index, percentage of albumen, yolk, and eggshell weight, eggshell thickness, yolk color score, Haugh unit (HU), albumen and yolk index) and malondialdehyde (MDA) levels in egg yolk. The cold-press extraction method offers advantages in terms of cost-effectiveness and environmental sustainability. This technique avoids the use of solvents and heat, which helps preserve thermosensitive compounds. However, a notable limitation is that the extracted compounds remain in their complex, non-degraded forms, which may limit the efficiency of isolating certain bioactive constituents—especially such as phenolics, flavonoids, and alkaloids that typically require solvents for effective extraction.

**Results:** Phytochemical screening conducted in this study revealed low concentrations of several compounds, and notably, the extract of *S. nigrum* fruit obtained using a cold-press juicer method showed no presence of triterpenoids or steroids. The results showed that black

## Keywords:

Antioxidant, Cold-press extraction, Egg quality

## \* Corresponding Author:

Widya Hermana, PhD.

Address: Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, IPB University, Bogor, Indonesia.

Tel: +62 (251) 8626213

E-mail: [widyaha@apps.ipb.ac.id](mailto:widyaha@apps.ipb.ac.id)

Copyright © 2026 The Author(s).  
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>).  
Noncommercial uses of the work are permitted, provided the original work is properly cited.

nightshade extract supplementation did not significantly affect quail performance ( $P>0.05$ ). However, significant improvements ( $P<0.05$ ) were observed in physical egg quality parameters, particularly in albumen index, yolk color, and HU at higher doses (T2 and T3). The MDA content in the yolk significantly decreased with increased extract concentration.

**Conclusion:** Under the conditions of this study, *S. nigrum* L. fruit extract can be considered a promising natural phytobiotic additive to enhance internal egg quality and oxidative stability without compromising laying performance in *C. coturnix japonica*.

## 1. Introduction

**C***oturnix coturnix japonica* is one of the poultry species widely raised in Indonesia. It has been domesticated and utilized by local communities as a source of both eggs and meat. Quail possess a relatively short reproductive cycle, requiring only about 45 days to mature. Annual egg production per bird can reach approximately 270–300 eggs. The average weight of a quail egg ranges from 11 to 13 grams, which is relatively small compared to chicken eggs and those of other poultry species. Despite their size, quail eggs are rich in nutrients and are considered an excellent dietary source of protein for humans [1].

One promising approach to enhance quail productivity is through the use of local plants as phytobiotics. Phytobiotics are plant-derived bioactive compounds that are environmentally friendly and capable of boosting both immunity and production performance in poultry [2]. Phytobiotics are considered a viable alternative to antibiotic growth promoters (AGPs). The antioxidant components found in phytobiotics help protect body cells from damage caused by free radicals and oxidative stress, and they may also stimulate the expression of endogenous antioxidant enzymes [3]. Plants that contain high levels of bioactive compounds have great potential for use as phytobiotics in animal feed.

Several studies have explored the application of phytobiotics in poultry. Krauze [4] reported that supplementing cinnamon oil at 0.25 mL/L in the drinking water of chickens improved performance, metabolism, immune response, and antioxidant activity. Another study demonstrated that administering 2 mL/L of *Bifidobacterium* sp. and *Guazuma ulmifolia* leaf extracts in quail drinking water increased egg mass, Haugh unit (HU), and yolk color intensity [5].

*Solanum nigrum*, commonly known as black nightshade or “leunca” in Indonesia, is a local plant known to contain various bioactive compounds with strong

antioxidant activity [6]. Given these properties, the use of *S. nigrum* fruit extract as a phytobiotic presents significant potential. Therefore, further studies are necessary to evaluate the effects of *S. nigrum* fruit extract as a phytobiotic in laying quail.

## 2. Materials and Methods

### 2.1. Experimental design

The experiment consisted of four treatment groups with four replicates each. Quail rearing was conducted at Arkan Quail Farm, located in Bogor, Indonesia. Evaluation of egg quality was performed at the Poultry Nutrition Laboratory, Department of Nutrition and Feed Technology, Faculty of Animal Science, IPB University. A total of 160 female Japanese quails (*C. coturnix japonica*) were used, reared from 6 to 16 weeks of age. The birds were provided with a commercial feed (New Hope P100, PT New Hope Farm Indonesia), and both feed and drinking water were supplied ad libitum. Feeding, watering, and cage cleaning were performed daily. Black nightshade (*S. nigrum*) fruit extract was administered every day. Eggs were collected and weighed daily for data recording. Physical egg quality assessment was conducted every two weeks, with three eggs per replication selected using a simple random sampling method. The parameters observed included productivity performance: feed intake (g/bird/day), quail-day production (QDP [%]), egg weight (g), egg mass (g/bird/day), feed conversion ratio (FCR), and water intake (mL/bird/day). Egg physical quality was evaluated, including yolk, albumen, and shell weight and percentage, egg index, yolk index, albumen index, shell thickness, yolk color, and HU. The study also measured malondialdehyde (MDA) content (mg/kg) of the eggs.

### 2.2. Black nightshade fruit extraction

The extraction method was a modified procedure based on the technique described by Puspitaningrum [7]. The fruits were thoroughly washed and drained pri-

or to extraction. The cleaned fruits were processed using a cold-press juicer to obtain the crude extract, which was subsequently filtered and pasteurized for 10 minutes. The resulting black nightshade fruit extract was stored in a refrigerator at 4 °C until further use.

### 2.3. Statistical analyses

Data were analyzed using analysis of variance (ANOVA) with IBM SPSS software, version 25. When significant differences were found ( $P < 0.05$ ), further comparisons between treatment means were conducted using Tukey's post hoc test. The MDA levels in egg yolks were analyzed descriptively. The study included four treatment groups: T0=drinking water (control), T1=1 L drinking water + 2 mL of black nightshade fruit extract, T2=1 L drinking water + 4 mL of black nightshade fruit extract, and T3=1 L drinking water + 6 mL of black nightshade fruit extract.

## 3. Results

### 3.1. Phytochemical analysis of fruit extraction

The results of the phytochemical screening of black nightshade (*S. nigrum*) fruit extract are presented in Table 1. The extract was found to contain alkaloids, saponins, tannins, phenolics, flavonoids, and glycosides. However, triterpenoids and steroids were not detected in the extract.

### 3.2. Production performance of laying quails aged 6–16 weeks

The production performance of laying quails aged 6 to 16 weeks supplemented with black nightshade (*S. nigrum*) fruit extract in drinking water is summarized in Table 2. The supplementation did not result in significant changes in feed intake, egg production, egg weight, egg mass, FCR, or water intake ( $P > 0.05$ ).

**Table 1.** Phytochemical screening of black nightshade fruit extract

Phytochemical	Result
Alkaloids	+
Saponins	+
Tanins	+
Phenolics	+
Flavonoids	+
Glycosides	+
Triterpenoids	-
Steroids	-

**Table 2.** Production performance of laying quails (6-16 week)

Variables	Mean±SD			
	Treatment			
	T0	T1	T2	T3
Feed intake (g/bird/day)	23.89±1.16	22.63±1.03	23.55±0.9	23.83±0.67
QDP (%)	83.42±11.95	78.83±9.78	80.45±14.22	84.87±12.63
Egg weight (g/egg)	11.2±0.23	10.85±0.32	11.07±0.22	11.06±0.32
Egg mass (g/bird/day)	9.24±1.44	8.52±1.09	8.93±1.71	9.40±1.5
FCR	2.63±0.58	2.73±0.67	2.84±1.19	2.63±0.7
Water intake (mL/bird/day)	67.68±11.25	64.28±3.77	64.87±7.03	61.62±9.99

**Table 3.** Egg quality of laying quails (6-16 week)

Variables	Treatment			
	T0	T1	T2	T3
Albumen weight (g)	7.76±0.63	7.63±0.6	7.89±0.5	7.92±0.57
Yolk weight (g)	3.70±0.43	3.64±0.53	3.72±0.35	3.67±0.39
Shell weight (g)	1.53±0.2	1.49±0.15	1.55±0.13	1.55±0.15
Albumen percentage (%)	67.75±2.27	67.75±3.4	67.98±2.05	68.37±2.29
Yolk percentage (%)	32.35±2.27	32.25±3.4	32.02±2.05	31.63±2.29
Shell percentage (%)	13.25±1.17	13.2±1.15	13.31±0.89	13.34±0.9
Egg index	78.16±2.86	78.41±2.83	78.16±2.43	78.5±1.95
Albumen index	11.11±2.82 <sup>a</sup>	11.98±1.74 <sup>ab</sup>	12.46±2.27 <sup>b</sup>	12.29±2.18 <sup>ab</sup>
Yolk index	44.35±2.95	43.63±3.07	43.17±3.46	43.32±2.31
Yolk colour	3.58±0.69 <sup>a</sup>	4.41±0.5 <sup>b</sup>	4.41±0.89 <sup>b</sup>	4.5±0.2 <sup>b</sup>
HU	92.67±1.34 <sup>a</sup>	94.16±0.55 <sup>ab</sup>	95.16±1.5 <sup>b</sup>	95.55±1.03 <sup>b</sup>
Shell thickness (mm)	0.2±0	0.19±0.01	0.2±0	0.2±0.01

Different letters on the same row indicate a statistical difference ( $P \leq 0.05$ ) among treatments.

### 3.3. Egg quality of laying quails aged 6–16 weeks

Egg quality parameters of laying quails from 6 to 16 weeks of age are shown in Table 3. Significant improvements were observed in albumen index, yolk color, and HU ( $P < 0.05$ ). Among the groups, T3 showed the highest values in these three parameters. No significant differences were found in egg index, yolk index, eggshell percentage, or shell thickness ( $P > 0.05$ ).

### 3.4. MDA content in eggs

The levels of MDA and the percentage reduction across treatment groups are presented in Table 4. The control group (T0) had an MDA level of 1.776 mg/kg. Treatment groups T2 and T3 showed reductions of 23.2% and

28.38%, respectively, with the T3 group exhibiting the greatest decrease in MDA concentration.

## 4. Discussion

Phytochemical screening conducted in this study revealed low concentrations of several compounds, and notably, the extract of *S. nigrum* fruit obtained using a cold-press juicer method showed no presence of triterpenoids or steroids. Hydroalcoholic extraction of *Solanum nigrum* fruit from India revealed the absence of alkaloids, while the ethyl acetate method yielded no alkaloids or terpenoids, though the flavonoid content reached 10.46% [8].

**Table 4.** Egg yolk MDA levels of 16-week-old laying quails

Treatment	MDA (mg/kg)	MDA Reduction (%)
T0	1.776	-
T1	1.89	-6.42
T2	1.364	23.2
T3	1.272	28.38

The discrepancies between the phytochemical findings of the present study and previous research may be attributed to differences in fruit origin and in the extraction methods. Different extraction techniques can yield distinct profiles of bioactive compounds. The cold-press extraction method offers advantages in terms of cost-effectiveness and environmental sustainability. This technique avoids the use of solvents and heat, which helps preserve thermosensitive compounds. However, a notable limitation is that the extracted compounds remain in their complex, non-degraded forms, which may limit the efficiency of isolating certain bioactive constituents—especially those such as phenolics, flavonoids, and alkaloids that typically require solvents for effective extraction [9].

The effects of *Solanum nigrum* fruit extract supplementation in quail drinking water on performance during the study are presented in Table 1. The average feed intake ranged from 22.63 to 23.89 g/bird/day. The extract treatments did not negatively affect feed consumption. Analysis of variance showed no significant differences ( $P>0.05$ ), likely due to the use of the same feed composition across all treatments, resulting in uniform energy intake.

Egg production did not differ significantly ( $P>0.05$ ) following black nightshade extract supplementation. Egg weight was also not significantly affected by treatment ( $P>0.05$ ). The average egg weight ranged from 10.85 to 11.20 g/egg. The absence of differences may be attributed to the uniform protein content in the feed provided to all groups. As reported by Ojediran [10], both egg production and egg weight are heavily influenced by protein intake and feed consumption levels. Egg mass was likewise unaffected ( $P>0.05$ ) by *S. nigrum* supplementation. The average egg mass ranged from 8.52 to 9.40 g/bird/day. Egg mass in the T3 group was slightly higher than that of T0. This increase may result from the combined effects of protein intake, egg weight, and egg production.

FCR showed no significant difference among treatments ( $P>0.05$ ). The FCR ranged from 2.63 to 2.84. A lower FCR indicates higher feed efficiency, meaning quails were more effective in converting feed into eggs [2]. Water intake ranged from 61.62 to 67.68 mL/bird/day and was not significantly different between treatments ( $P>0.05$ ). Oxidative stress may influence water consumption, with higher intake possibly indicating elevated stress levels due to non-ideal temperatures [11]. During this study, birds were maintained at temperatures ranging from 24 to 36 °C, which may not represent optimal conditions.

The physical quality of eggs is shown in Table 2. The weight of albumen, yolk, and shell ranged from 7.63 to 7.92 g, 3.64 to 3.72 g, and 1.49 to 1.55 g, respectively. Egg white weight is generally proportional to total egg weight, which explains the relatively lower yolk weight in heavier eggs [12]. There were no significant differences in component percentages ( $P>0.05$ ), with albumen, yolk, and shell percentages ranging from 67.75% to 68.37%, 31.63% to 32.35%, and 13.2% to 13.34%, respectively. Compared to Chimezie [12], the present study yielded higher values for albumen and shell, while yolk proportions were similar.

HU values ranged from 92.67 to 95.55 and showed a significant difference ( $P<0.05$ ). These values are consistent with Lovela [5] and higher than those in Chimezie [12]. HU is influenced by albumen height and protein quality. Flavonoids present in the extract may contribute positively by enhancing nutrient absorption and improving protein metabolism via proteolytic enzyme activity [13, 14]. Egg and yolk indices showed no significant differences ( $P>0.05$ ), while the albumen index was significantly affected ( $P<0.05$ ). Indices observed were 78.16–78.50 (egg), 11.11–12.46 (albumen), and 43.17–44.35 (yolk). The albumen index correlates with HU and albumen height [15].

Yolk color was significantly affected by treatment ( $P<0.05$ ), ranging from 3.58 to 4.50, although lower than those reported by Lovela [5]. The color enhancement is likely due to carotenoid content in *S. nigrum*, particularly  $\beta$ -carotene [16, 17]. Shell thickness showed no significant difference ( $P>0.05$ ), ranging from 0.19 to 0.20 mm. A thicker shell offers better protection against microbial contamination and water loss [18, 19]. Yolk MDA levels ranged from 1.272 to 1.890 mg/kg. MDA is a marker of oxidative stress resulting from lipid peroxidation. T2 and T3 treatments reduced MDA levels by 23.2% and 28.38%, respectively, compared to T0, suggesting antioxidant effects of *S. nigrum*. The extract contains flavonoids, carotenoids, phenols, tannins, and alkaloids, which may contribute to this reduction [16, 20].

## 5. Conclusion

Supplementation of *S. nigrum* L. fruit extract in the drinking water of laying quails (*C. coturnix japonica*) did not significantly affect production performance ( $P>0.05$ ), but improved selected internal egg quality parameters, particularly albumen index, yolk color, and HU ( $P<0.05$ ). Yolk MDA levels were reduced in the higher supplementation groups, indicating decreased lipid peroxidation. Under the conditions of the present

study, *S. nigrum* fruit extract shows promise as a natural phyto-biotic additive to enhance internal egg quality and oxidative stability without adverse effects on laying performance.

### Acknowledgements

The author would like to thank the lecturers and technicians of the Poultry Nutrition Division and the Arkan Quail Farm team who helped to perform this study.

### Compliance with ethical guidelines

All animal procedures were approved by the Animal Ethics Committee, School of Veterinary Medicine and Biomedical Science, **IPB University**, Bogor, Indonesia (Code: 274/KEH/SKE/XIII/2024).

### Data availability

The data that support the findings of this study are available upon request from the corresponding author.

### Funding

This research did not receive any grant from funding agencies in the public, commercial, or non-profit sectors.

### Authors' contributions

Conceptualization, study design, experiments, data interpretation, project administration, technical, and material support: All authors; Data acquisition, statistical analysis, and writing: Syauqi Syah Dewan.

### Conflict of interest

The authors declared no conflict of interest.

### References

- [1] Saraswati TR, Tana S. Development of Japanese quail (*Coturnix coturnix japonica*) embryo. *Int J Sci Eng*. 2015; 8(1):38-41. [DOI:10.12777/ijse.8.1.38-41]
- [2] Ampode KM. Papaya (*Carica papaya* Linn.) leaf meal as phyto-biotic feed supplement improves the production performance of growing Japanese quails. *Int J Biosci*. 2019; 14(5):534-41. [Link]
- [3] Kikusato M. Phyto-biotics to improve health and production of broiler chickens: functions beyond the antioxidant activity. *Anim Biosci*. 2021; 34(3):345-53. [DOI:10.5713/ab.20.0842] [PMID]
- [4] Krauze M, Cendrowska-Pinkosz M, Matusevičius P, Stępniewska A, Jurczak P, Ognik K. The effect of administration of a phyto-biotic containing cinnamon oil and citric acid on the metabolism, immunity, and growth performance of broiler chickens. *Animals*. 2021; 11(2):399. [DOI:10.3390/ani11020399] [PMID]
- [5] Lovela AR, Lokapirnasari WP, Al-Arif MA, Soeharsono S, Hidanah S, Warsito SH, et al. The quality of Japanese quail eggs after administration of *Bifidobacterium* sp. and *Guzuma ulmifolia* Leaf Extract. *J Med Vet*. 2023; 6(1):132-6. [DOI:10.20473/jmv.vol6.iss1.2023.132-136]
- [6] Alam A, Sahar A, Sameen A, Faisal MN. The effects of bioactive components in *Solanum nigrum* against oxidative stress in liver damage. *Food Sci Technol*. 2022; 42:e61822. [DOI:10.1590/fst.61822]
- [7] Puspitaningrum T, Mahfudz LD, Nasoetion MH. [The potency of garlic (*Allium sativum*) and *Lactobacillus acidophilus* as syn-biotics on broiler performance (Indonesian)]. *J Sain Peternak Indones*. 2021; 16(2):210-4. [DOI:10.31186/jspi.id.16.2.210-214]
- [8] Goel K, Singh R, Saini V, Bansal S. Phyto-chemical estimation and evaluation of antioxidant activity of hydro-alcoholic and ethylacetate extracts of *Tribulus terrestris* linn. (*gokhru*) and *Solanum nigrum* linn (*makoi*). *Int J Health Sci*. 2022; 6(56):7661-76. [DOI:10.53730/ijhs.v6n56.12158]
- [9] Al Juhaimi F, Özcan MM, Ghafoor K, Babiker EE, Hussain S. Comparison of cold-pressing and soxhlet extraction systems for bioactive compounds, antioxidant properties, polyphenols, fatty acids and tocopherols in eight nut oils. *J Food Sci Technol*. 2018; 55(8):3163-73. [DOI:10.1007/s13197-018-3244-5] [PMID]
- [10] Ojediran T, Ganiyu B, Muhammed A, Emiola A. Egg production, economic indices, and external and internal egg quality parameters of laying Japanese quails (*Coturnix japonica*) fed palm (*Elaeis guineensis*) kernel cake. *Anim Sci Genet*. 2022; 18(2):1-11. [DOI:10.5604/01.3001.0015.8489]
- [11] Fadhila R, Ulupi N, Maheshwari H. Utilization of coconut shell liquid smoke on quail performance at grower period. *J Ilmu Prod dan Tek Hasil Peternakan*. 2023; 11(3):163-9. [DOI:10.29244/jipthp.11.3.163-169]
- [12] Chimezie VO, Egbeyan Ahmed J, Faseun T. Effect of *Phyllanthus amarus* supplemented diet on egg production and egg quality traits of Japanese quail. *Int J Anim Sci Husb Livest Prod*. 2022; 8(1):423-9. [Link]
- [13] Deko MK, Djunaidi IH, Halim Natsir MH. The use of garlic (*Allium sativum* Linn) bulb and husk powder as an organic feed additive on egg quality of laying hens. *Int Res J Adv Eng Sci*. 2018; 3(3):218-22. [Link]
- [14] Salazar I, Rodriguez R, Aroche R, Valdivié M, Martínez Y. Phyto-biotic effect of *Jatropha curcas* leaf powder on productivity, egg quality and blood biochemistry of laying quails. *Cuba J Agric Sci*. 2021; 55(3):315-26. [Link]
- [15] Al-Hadeedy IY, Mohammed MA, Abdulazeez ST, Ameen QA, Shaker AS. Genetic correlation of exterior and interior quality characteristics in white quail egg. *J Surv Fish Sci*. 2023; 10(3S):949-59. [Link]

- [16] Khan HJ, Ahmad MK, Khan AR, Rastogi N, Mahdi AA, Ansari JA, et al. Identification of anticancer and antioxidant phytoconstituents from chloroform fraction of *Solanum nigrum* L. Berries using GC-MS/MS analysis. *Indian J Exp Biol.* 2016; 54(11):774-82. [\[Link\]](#)
- [17] Jithendran L, Kalpana C. Nutritional potential of *Solanum nigrum* linn berries grown in home garden. *Int J Food Nutr Sci.* 2021; 10(1):16. [\[Link\]](#)
- [18] Kocetkovs V, Radenkovs V, Juhnevica-Radenkova K, Jakovlevs D, Muizniece-Brasava S. The impact of eggshell thickness on the qualitative characteristics of stored eggs produced by three breeds of laying hens of the cage and cage-free housed systems. *Appl Sci.* 2022; 12(22):11539. [\[DOI:10.3390/app122211539\]](#)
- [19] Liu Y, Song M, Bai H, Wang C, Wang F, Yuan Q. Curcumin improves the egg quality, antioxidant activity, and intestinal microbiota of quails during the late laying period. *Poult Sci.* 2024; 103(1):103233. [\[DOI:10.1016/j.psj.2023.103233\]](#) [\[PMID\]](#)
- [20] Meerwal P, Jain GC. Effect of *Solanum nigrum* L. Fruit extract on serum sex hormones and testis of male wistar rats. *Int J Sci Technol Res.* 2019; 8(12):3496-502. [\[Link\]](#)

This Page Intentionally Left Blank